

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 January 2001 (25.01.2001)

PCT

(10) International Publication Number
WO 01/05828 A1

(51) International Patent Classification⁷: **C07K 14/47**,
16/18, G01N 33/53, 33/567, C12N 5/10, 15/12, 15/63,
15/64

0907 (US). SWANSON, Richard [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). LIU, Yuan [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). LAGRUTTA, Armando [PA/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(21) International Application Number: PCT/US00/19585

(22) International Filing Date: 18 July 2000 (18.07.2000)

(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(25) Filing Language: English

(81) Designated States (*national*): CA, JP, US.

(26) Publication Language: English

(84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

(30) Priority Data:
60/144,764 20 July 1999 (20.07.1999) US

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

(71) Applicant (*for all designated States except US*): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): UEBELE, Victor [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/05828 A1

(54) Title: NOVEL HUMAN CALCIUM SENSITIVE POTASSIUM CHANNEL SUBUNITS

(57) Abstract: The present invention is directed to novel human DNA sequences encoding calcium sensitive potassium channel subunits $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$, the proteins encoded by the DNA sequences, vectors comprising the DNA sequences, host cells containing the vectors, and methods of identifying inhibitors and agonists of calcium sensitive potassium channels containing human $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits and inhibitors and agonists of $\beta 3$ gene transcription.

TITLE OF THE INVENTION

NOVEL HUMAN CALCIUM SENSITIVE POTASSIUM CHANNEL SUBUNITS

CROSS-REFERENCE TO RELATED APPLICATIONS

5 Not applicable.

STATEMENT REGARDING FEDERALLY-SPONSORED R&D

Not applicable.

10 REFERENCE TO MICROFICHE APPENDIX

Not applicable.

FIELD OF THE INVENTION

15 The present invention is directed to novel human DNA sequences
encoding subunits of calcium sensitive potassium channels.

BACKGROUND OF THE INVENTION

Voltage-gated potassium channels form transmembrane pores that open or close in response to changes in cell membrane potential and selectively allow potassium ions to pass through the membrane. Voltage-gated potassium channels have been found in cells traditionally considered both excitable (*e.g.*, neurons, myocytes, secretory cells) and non-excitable (*e.g.*, T-cells, osteoclasts) and have been shown to maintain cell membrane potential and control the repolarization of action potentials in such cells. Following depolarization, voltage-gated potassium channels open, allowing potassium efflux and thus membrane repolarization. This behavior has made voltage-gated potassium channels important targets for drug discovery in connection with a variety of diseases. As a result, many voltage-gated potassium channels have been identified and many cloned. They are distinguishable by differences in primary structure and tissue-specific patterns of expression, as well as by electrophysiological and pharmacological properties. For reviews of voltage-gated potassium channels see Robertson, 1997, Trends Pharmacol. Sci. 18:474-483; Jan & Jan, 1997, J. Physiol. 505:267-282; Catterall, 1995, Ann. Rev. Biochem. 64:493-531.

20
25
30

Many functional voltage-gated potassium channels are believed to be tetramers of four α subunits, each of which contains six transmembrane spanning

segments. The α subunits making up a tetramer may be the same (in the case of homotetramers) or may be different (in the case of heterotetramers). The membrane-spanning α subunits making up the tetramers may sometimes be associated with additional, β subunits, which may alter the behavior of the α subunits.

- 5 A particular type of voltage-gated potassium channel is the voltage-gated and calcium sensitive potassium channel, also known as the calcium sensitive potassium channel. Calcium sensitive potassium channels are present in a wide variety of cells and are unique among voltage-gated potassium channels because their activity is regulated not only by changes in membrane potential but also by
- 10 intracellular calcium concentration. Plasma membrane depolarization and increases in cytoplasmic calcium concentration both raise the open probability of calcium sensitive potassium channels. Therefore, calcium sensitive potassium channels can serve as a link between cellular processes involving increases in intracellular calcium and membrane excitability. Calcium sensitive potassium channels are believed to
- 15 play a negative feedback role by terminating signaling events involving an increase in intracellular calcium, *e.g.*, glucose mediated insulin release, blood vessel muscle tone, bronchial airway smooth muscle tone, and regulation of intraocular pressure. (Tanaka et al., 1997, J. Physiol. 502:545-557; Kaczorowski et al., 1996, J. Bioenerg. Biomem. 28:255-267; Vergara et al., 1998, Curr. Opin. Neurobiol. 8:321-329).
- 20 Certain calcium sensitive potassium channels have been isolated and studied. Functional calcium sensitive potassium channels are composed of α subunits that may be associated with smaller β subunits. The α subunit is believed to form the channel pore while a previously described β subunit increases the calcium sensitivity of the channel and makes the channel susceptible to regulation by certain substances,
- 25 *e.g.*, dehydrosoyasaponin (McManus et al., 1995, Neuron 14:645-650). The calcium sensitive potassium channel from bovine tracheal smooth muscle was purified and shown to be composed of an ~130 kDa α subunit and a 31 kDa β subunit (Garcia-Calvo et al., 1994, J. Biol. Chem. 269:676-682). Tseng-Crank et al. (1994, Neuron 13:1315-1330) cloned nine related calcium sensitive potassium channel α subunits
- 30 from human brain. These α subunits are thought to be splice variants derived from a single gene, the *h-slo* gene (Tseng-Crank et al., 1994, Neuron 13:1315-1330). Knauss et al., 1994, J. Biol. Chem. 269:17274-17278 purified and cloned a β subunit of a calcium sensitive potassium channel from tracheal smooth muscle.

In most cells, the opening of calcium sensitive potassium channels results in the generation of non-inactivating, hyperpolarizing potassium currents. However, in certain cells (*e.g.*, chromaffin cells of the adrenal gland and hippocampal neurons), the currents are inactivating. Following the discovery of the invention described herein, Wallner et al., 1999, Proc. Natl. Acad. Sci. USA 96:4137-4132 disclosed the existence of the human $\beta 2$ calcium sensitive potassium channel subunit that, when combined with the α subunit, formed inactivating calcium sensitive potassium channels. The ability to confer inactivation was ascribed to the N-terminal 19 amino acids of the $\beta 2$ subunit.

U.S. Patent No. 5,776,734 is directed to nucleic acids encoding the bovine and human $\beta 1$ subunit of the calcium sensitive potassium channel. U.S. Patent No. 5,637,470 is directed to methods of identifying compounds that modulate the activity of calcium sensitive potassium channels.

SUMMARY OF THE INVENTION

The present invention is directed to novel human DNA sequences encoding β subunits of calcium sensitive potassium channels. The present invention includes DNAs that encode the β subunits $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ of human calcium sensitive potassium channels. The DNAs comprise the nucleotide sequences shown in SEQ.ID.NO.:1 ($\beta 2$), SEQ.ID.NO.:3 ($\beta 3a$), SEQ.ID.NO.:5 ($\beta 3b$), SEQ.ID.NO.:7 ($\beta 3c$), and SEQ.ID.NO.:9 ($\beta 3d$). Also provided are proteins encoded by the novel DNA sequences. The proteins comprise the deduced amino acid sequences shown in SEQ.ID.NO.:2 ($\beta 2$), SEQ.ID.NO.:4 ($\beta 3a$), SEQ.ID.NO.:6 ($\beta 3b$), SEQ.ID.NO.:8 ($\beta 3c$), and SEQ.ID.NO.:10 ($\beta 3d$). Methods of expressing the novel subunit proteins in recombinant systems are provided as well as methods of identifying activators and inhibitors of potassium channels comprising the subunits.

The present invention also includes a genomic DNA fragment containing the 5' portions of the $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ subunits, as well as the 5' portion of the core portion of the $\beta 3$ subunits. This genomic DNA fragment contains promoter elements for the subunits. Methods of screening for compounds which affect transcription of the gene encoding the $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ subunits are also provided.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A shows a DNA sequence encoding the $\beta 2$ subunit of the human calcium sensitive potassium channel (SEQ.ID.NO.:1). The start ATG codon is at position 271-273; the stop codon is at position 976-978. Figure 1B shows the deduced amino acid sequence (SEQ.ID.NO.:2) of the $\beta 2$ subunit.

Figure 2A shows a DNA sequence encoding the $\beta 3a$ subunit of a human calcium sensitive potassium channel (SEQ.ID.NO.:3). The start ATG codon is at position 341-343; the stop codon is at position 1172-1174. Figure 2B shows the deduced amino acid sequence (SEQ.ID.NO.:4) of the $\beta 3a$ subunit.

Figure 3A shows a DNA sequence encoding the $\beta 3b$ subunit of a human calcium sensitive potassium channel (SEQ.ID.NO.:5). The start ATG codon is at position 796-798; the stop codon is at position 1567-1569. Figure 3B shows the deduced amino acid sequence (SEQ.ID.NO.:6) of the $\beta 3b$ subunit.

Figure 4A shows a DNA sequence encoding the $\beta 3c$ subunit of a human calcium sensitive potassium channel (SEQ.ID.NO.:7). The start ATG codon is at position 869-871; the stop codon is at position 1694-1696. Figure 4B shows the deduced amino acid sequence (SEQ.ID.NO.:8) of the $\beta 3c$ subunit.

Figure 5A shows a DNA sequence encoding the $\beta 3d$ subunit of a human and calcium sensitive potassium channel (SEQ.ID.NO.:9). The start ATG codon is at position 457-459; the stop codon is at position 1294-1296. Figure 5B shows the deduced amino acid sequence (SEQ.ID.NO.:10) of the $\beta 3d$ subunit.

Figure 6 shows an alignment of the deduced amino acid sequences of the human calcium sensitive potassium channel $\beta 1$ (SEQ.ID.NO.:11), $\beta 2$ (SEQ.ID.NO.:2), $\beta 3a$ (SEQ.ID.NO.:4), $\beta 3b$ (SEQ.ID.NO.:6), $\beta 3c$ (SEQ.ID.NO.:8), and $\beta 3d$ (SEQ.ID.NO.:10) subunits.

Figure 7 shows the effect of the co-expression of the novel β subunits of the present invention on the electrophysiological properties of the ion channel formed by the α subunit of a human calcium sensitive potassium channel. Figure 7A shows the current-voltage relations recorded in inside-out patches expressing calcium sensitive potassium channel α or α and β subunits. α and β subunit cRNAs were co-injected in 1:10 molar ratio (β in excess) to detect maximum effects. The voltage clamp protocol consisted of a pre-pulse to -160 mV (200 ms), followed by 20 mV depolarizing steps from -80 to +80 mV (500 ms); holding potential was -80 mV; internal Ca^{2+} was 30 μM . Subunits $\beta 3b$ and $\beta 3d$ did not induce noticeable changes in

the kinetics and voltage dependence of the channels formed by α subunits, although they might decrease current density. Figure 7B: Boltzmann equations were fit to normalized conductances for the records shown in 7A, which were calculated from peak currents and plotted as function of test potential. $V_{1/2}$ values are: 20 mV (5 α subunit alone); -55 mV (α + $\beta 2$ subunit); 45.36 mV (α + $\beta 3a$ subunit); 20 mV (α + $\beta 3c$ subunit). Figure 7C shows that co-expression of $\beta 3$ subunit RNAs in molar excess of α subunit RNAs (up to 10X) reduced, but did not eliminate, a non-inactivating component of calcium sensitive potassium channel current. Inactivation rates and fractional inactivating current were calculated as described in Example 2.

10 Figure 8A-N shows the genomic sequence of GenBank accession number AC007823.4 (SEQ.ID.NO.:20). The different splice variants of the $\beta 3$ subunits are contained in nucleotides 1-40,467. The $\beta 3a$ -specific sequence is at positions 17,404-17,806; the $\beta 3b$ -specific sequence is at positions 24,710-25,507; the $\beta 3c/d$ sequence is at positions 32,590-33,514; the beginning of the $\beta 3$ core sequence

15 is at positions 33,515-33,705. The sequences involved in tissue specific expression (*e.g.*, promoters, enhancers, repressors) are likely to be located in nucleotides 1-17,404.

DETAILED DESCRIPTION OF THE INVENTION

20 For the purposes of this invention:

"Substantially free from other proteins" means at least 90%, preferably 95%, more preferably 99%, and even more preferably 99.9%, free of other proteins. Thus, a human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein preparation that is substantially free from other proteins will contain, as a

25 percent of its total protein, no more than 10%, preferably no more than 5%, more preferably no more than 1%, and even more preferably no more than 0.1%, of non-human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Whether a given human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein preparation is substantially free from other proteins can be

30 determined by conventional techniques of assessing protein purity such as, *e.g.*, sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) combined with appropriate detection methods, *e.g.*, silver staining or immunoblotting.

"Substantially free from other nucleic acids" means at least 90%, preferably 95%, more preferably 99%, and even more preferably 99.9%, free of other

nucleic acids. Thus, a human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit DNA preparation that is substantially free from other nucleic acids will contain, as a percent of its total nucleic acid, no more than 10%, preferably no more than 5%, more preferably no more than 1%, and even more preferably no more than 0.1%, of non-human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit nucleic acids. Whether a given human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit DNA preparation is substantially free from other nucleic acids can be determined by conventional techniques of assessing nucleic acid purity such as, *e.g.*, agarose gel electrophoresis combined with appropriate staining methods, *e.g.*, ethidium bromide staining, Northern or Southern blotting, or by sequencing.

A "conservative amino acid substitution" refers to the replacement of one amino acid residue by another, chemically similar, amino acid residue. Examples of such conservative substitutions are: substitution of one hydrophobic residue (isoleucine, leucine, valine, or methionine) for another; substitution of one polar residue for another polar residue of the same charge (*e.g.*, arginine for lysine; glutamic acid for aspartic acid); substitution of one aromatic amino acid (tryptophan, tyrosine, or phenylalanine) for another.

A polypeptide has "substantially the same biological activity as human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit" if that polypeptide is able to combine with a human calcium sensitive potassium channel α subunit thereby forming a functional potassium channel where the polypeptide confers upon the α subunit properties similar to those conferred by the $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits and where the polypeptide has an amino acid sequence that is at least about 50% identical to SEQ.ID.NO.:2, 4, 6, 8, or 10 when measured by such standard programs as BLAST or FASTA. For example, a polypeptide that is 50% identical in amino acid sequence to $\beta 3a$ (SEQ.ID.NO.:4) and is able to confer upon the α subunit properties such that electrophysiological measurements of the ion channel formed by the polypeptide and the α subunit result in graphs such as those shown in Figure 7A-C for the $\beta 3a$ subunit and the α subunit is a polypeptide that has "substantially the same biological activity as human calcium sensitive potassium channel $\beta 3a$ subunit."

The present invention relates to the identification and cloning of DNAs encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$

subunits, components of human calcium sensitive potassium channels. Expressed sequence tags (ESTs) (GenBank accession numbers AA904191, AI299145 and AI301175) were identified by searching databases for sequences with homology to the $\beta 1$ subunit. The cDNAs encoding the ESTs were purchased and sequenced in both
5 directions. The clone encoding AA904191 was determined to encode the entire $\beta 2$ subunit, since it contained in frame stop codons 5' to the start ATG of the open reading frame and the entire open reading frame.

The $\beta 2$ coding sequence was then used to search the databases for additional β subunits. Contigs were assembled from the identified ESTs and used to
10 search the database once again. Several ESTs were identified in this iterative manner (GenBank accession numbers AA195381, AA236930, AA236968, AA279911, AA761761 and AA934876). Available cDNAs encoding these ESTs were purchased and sequenced in both directions. None of these clones were full length. Because most were isolated in a preparation of tonsils enriched for B-cells, we performed 5'
15 RACE (rapid amplification of cDNA ends) using gene-specific oligonucleotides in the 3' untranslated region (UTR) and commercially prepared cDNA from human spleen, another tissue rich in B-cells (Clontech catalog # 7412-1), as the template. Multiple DNA fragments were amplified in this manner, cloned and sequenced in both directions. Sequencing revealed 4 subfamilies of full length clones, differing
20 only in their 5' ends: $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$.

The human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ subunits of the present invention exhibit tissue specific patterns of expression. Northern blotting of mRNAs isolated from various tissues has shown that the $\beta 2$ subunit is expressed predominately in uterus, heart, ovary, thyroid, fetal
25 kidney, adrenal medulla, and pancreas; the $\beta 3a$ subunit is expressed predominately in heart and skeletal muscle; the $\beta 3b$ subunit is expressed in most tissues examined except for brain, skeletal muscle and testes. The $\beta 3c$ and/or $\beta 3d$ subunits have been found in pancreas.

The tissue specific expression patterns of the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ subunits support the hypothesis that
30 these different subunits may contribute to the functional diversity of calcium sensitive potassium channels observed in different tissues. Activators and inhibitors of specific calcium sensitive potassium channels containing specific subunits may, therefore, have pharmacological efficacy in different pathological conditions, depending on the

subunit composition of the calcium sensitive potassium channels involved in the specific pathological condition.

Chromosomal mapping studies have shown that both the $\beta 2$ and $\beta 3$ subunits map to human chromosome 3q23-ter. The β subunits of the present invention have about 30-45% amino acid sequence identity to the previously known human $\beta 1$ subunit (GenBank accession no. U25138). The $\beta 2$ and $\beta 3$ subunits of the present invention have about 40% amino acid sequence identity to each other. The $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits differ only in their extreme N-terminal 1-20 amino acids, and are alternatively spliced variants of a single gene. Indeed, a genomic fragment of human DNA has been identified in the GenBank database that contains the 5' domains of $\beta 3a$, $\beta 3b$, $\beta 3c/d$, and the beginning of the conserved core in a contiguous fragment (accession number AC007823.4). See Figure 8. Additionally, two bacterial artificial chromosomes (BACs) have been isolated which contain the conserved core domain. One of these BACs also contains $\beta 3c/d$ specific sequence. Therefore, we have identified overlapping BAC clones that together encode the entire $\beta 3$ open reading frame. The $\beta 2$ subunit is encoded by a separate gene.

The present invention provides DNAs encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits that are substantially free from other nucleic acids. The present invention also provides isolated and/or recombinant DNA molecules encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits. The present invention provides DNA molecules substantially free from other nucleic acids comprising the nucleotide sequences shown in SEQ.ID.NO.:1, 3, 5, 7, or 9. cDNAs encoding each $\beta 3$ subunit have been isolated exhibiting a sequence polymorphism, encoding either a serine or an asparagine at the amino acid position that is equivalent to position 143 of $\beta 3b$. This represents amino acid 142 of the conserved core domain.

Accordingly, the present invention includes DNA substantially free from other nucleic acids as well as isolated and/or recombinant DNA encoding a polypeptide selected from the group consisting of: SEQ.ID.NO.:4; SEQ.ID.NO.:4 with an asparagine at position 163 instead of a serine; SEQ.ID.NO.:6; SEQ.ID.NO.:6 with a serine at position 143 instead of an asparagine; SEQ.ID.NO.:8; SEQ.ID.NO.:8 with an asparagine at position 161 instead of a serine; SEQ.ID.NO.:10; and SEQ.ID.NO.:10 with a serine at position 165 instead of an asparagine.

The present invention includes DNA substantially free from other nucleic acids as well as isolated and/or recombinant DNA encoding a polypeptide comprising the conserved $\beta 3$ core amino acid sequence, positions 2-246 of SEQ.ID.NO.:6.

5 The present invention includes isolated DNA molecules as well as DNA molecules that are substantially free from other nucleic acids comprising the coding regions of SEQ.ID.NOs.:1, 3, 5, 7, and 9. Accordingly, the present invention includes isolated DNA molecules and DNA molecules substantially free from other nucleic acids having a sequence comprising positions 271 to 975 of SEQ.ID.NO.:1,
10 positions 341 to 1171 of SEQ.ID.NO.:3, positions 796 to 1566 of SEQ.ID.NO.:5, positions 869 to 1693 of SEQ.ID.NO.:7, or positions 457 to 1293 of SEQ.ID.NO.:9.

 Also included are recombinant DNA molecules having a nucleotide sequence comprising positions 271-975 of SEQ.ID.NO.:1, positions 341 to 1171 of SEQ.ID.NO.:3, positions 796 to 1566 of SEQ.ID.NO.:5, positions 869 to 1693 of
15 SEQ.ID.NO.:7, or positions 457 to 1293 of SEQ.ID.NO.:9. The novel DNA sequences of the present invention encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits, in whole or in part, can be linked with other DNA sequences, *i.e.*, DNA sequences to which human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits are not naturally linked, to
20 form "recombinant DNA molecules" encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits. Such other sequences can include DNA sequences that control transcription or translation such as, *e.g.*, translation initiation sequences, internal ribosome entry sites, promoters for RNA polymerase II, transcription or translation termination sequences, enhancer sequences, sequences that
25 control replication in microorganisms, sequences that confer antibiotic resistance, or sequences that encode a polypeptide "tag" such as, *e.g.*, a polyhistidine tract, the FLAG epitope, the myc epitope, GST, or maltose binding protein. The novel DNA sequences of the present invention can be inserted into vectors such as plasmids, cosmids, viral vectors, P1 artificial chromosomes, or yeast artificial chromosomes.

30 The present invention also includes DNA substantially free from other nucleic acids as well as isolated and/or recombinant DNA comprising genomic sequences of the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits. The present invention includes DNA substantially free from other nucleic acids as well as isolated and/or recombinant DNA comprising

SEQ.ID.NO.:20; positions 1-40,467 of SEQ.ID.NO.:20; positions 17,404-17,806 of SEQ.ID.NO.:20; positions 24,710-25,507 of SEQ.ID.NO.:20; positions 32,590-33,514 of SEQ.ID.NO.:20; positions 33,515-33,705 of SEQ.ID.NO.:20; or positions 1-17,404 of SEQ.ID.NO.:20.

5 Included in the present invention are DNA sequences that hybridize to at least one of SEQ.ID.NOs:1, 3, 5, 7, 9, or 20 under conditions of high stringency. By way of example, and not limitation, a procedure using conditions of high stringency is as follows: Prehybridization of filters containing DNA is carried out for 2 hr. to overnight at 65°C in buffer composed of 5X SSC, 10X Denhardt's solution,
10 50% Formamide, 2% SDS and 100 µg/ml denatured salmon sperm DNA. Hybridization of 32P-labelled, random primed probe is carried out in 5X SSPE, 10X Denhardt's solution, 50% Formamide, 2% SDS, 100ug/ml salmon sperm DNA at 42°C overnight. Washing of filters is done in 2X SSC, 0.05% SDS at 42°C for 40 minutes, followed by 0.1X SSC, 0.05% SDS at 65°C for 40 minutes.

15 Other procedures using conditions of high stringency would include either a hybridization carried out in 5XSSC, 5X Denhardt's solution, 50% formamide at 42°C for 12 to 48 hours or a washing step carried out in 0.2X SSPE, 0.2% SDS at 65°C for 30 to 60 minutes.

20 Reagents mentioned in the foregoing procedures for carrying out high stringency hybridization are well known in the art. Details of the composition of these reagents can be found in, *e.g.*, Sambrook, Fritsch, and Maniatis, 1989, Molecular Cloning: A Laboratory Manual, second edition, Cold Spring Harbor Laboratory Press. In addition to the foregoing, other conditions of high stringency which may be used are well known in the art.

25 The degeneracy of the genetic code is such that, for all but two amino acids, more than a single codon encodes a particular amino acid. This allows for the construction of synthetic DNA that encodes the human calcium sensitive potassium channel β2, β3a, β3b, β3c, or β3d subunit proteins where the nucleotide sequence of the synthetic DNA differs significantly from the nucleotide sequences of
30 SEQ.ID.NOs:1, 3, 5, 7, or 9 but still encodes the same human calcium sensitive potassium channel β2, β3a, β3b, β3c, or β3d subunit proteins as SEQ.ID.NOs:2, 4, 6, 8, or 10. Such synthetic DNAs are intended to be within the scope of the present invention.

Mutated forms of SEQ.ID.NOs:1, 3, 5, 7, or 9 are intended to be within the scope of the present invention. In particular, mutated forms of SEQ.ID.NOs:1, 3, 5, 7, or 9 which encode proteins that either do not interact with an α subunit or which when combined with α subunits give rise to calcium sensitive potassium channels having altered voltage dependence, calcium sensitivity, current kinetics (such as activation, inactivation or deactivation), or pharmacologic properties as compared to wild-type calcium sensitive potassium channels are within the scope of the present invention. Such mutant forms can differ from SEQ.ID.NOs:1, 3, 5, 7, or 9 by having nucleotide deletions, substitutions, or additions.

Also intended to be within the scope of the present invention are RNA molecules having sequences corresponding to SEQ.ID.NOs:1, 3, 5, 7, or 9. Antisense nucleotides, DNA or RNA, that are the reverse complements of SEQ.ID.NOs:1, 3, 5, 7, or 9, or portions thereof, are also within the scope of the present invention. In addition, polynucleotides based on SEQ.ID.NOs:1, 3, 5, 7, or 9 in which a small number of positions are substituted with non-natural or modified nucleotides such as inosine, methyl-cytosine, or deaza-guanosine are intended to be within the scope of the present invention. Polynucleotides of the present invention can also include sequences based on SEQ.ID.NOs:1, 3, 5, 7, or 9 but in which non-natural linkages between the nucleotides are present. Such non-natural linkages can be, *e.g.*, methylphosphonates, phosphorothioates, phosphorodithionates, phosphoroamidites, and phosphate esters. Polynucleotides of the present invention can also include sequences based on SEQ.ID.NOs:1, 3, 5, 7, or 9 but having de-phospho linkages as bridges between nucleotides, *e.g.*, siloxane, carbonate, carboxymethyl ester, acetamidate, carbamate, and thioether bridges. Other internucleotide linkages that can be present include N-vinyl, methacryloxyethyl, methacrylamide, or ethyleneimine linkages. Peptide nucleic acids based upon SEQ.ID.NOs:1, 3, 5, 7, or 9 are also included in the present invention.

Another aspect of the present invention includes host cells that have been engineered to contain and/or express DNA sequences encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Such recombinant host cells can be cultured under suitable conditions to produce human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Such recombinant host cells are also useful in the methods of identifying activators and inhibitors of calcium sensitive potassium channels described herein. An

expression vector containing DNA encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can be used for the expression of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in a recombinant host cell. Recombinant host cells may be prokaryotic or eukaryotic, including but not limited to, bacteria such as *E. coli*, fungal cells such as yeast, mammalian cells including, but not limited to, cell lines of human, bovine, porcine, monkey and rodent origin, amphibian cells such as *Xenopus* oocytes, and insect cells including but not limited to *Drosophila* and silkworm derived cell lines. Cells and cell lines which are suitable for recombinant expression of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins and which are widely available, include but are not limited to, L cells L-M(TK⁻) (ATCC CCL 1.3), L cells L-M (ATCC CCL 1.2), 293 (ATCC CRL 1573), Raji (ATCC CCL 86), CV-1 (ATCC CCL 70), COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCC CCL 2), C127I (ATCC CRL 1616), BS-C-1 (ATCC CCL 26), MRC-5 (ATCC CCL 171), CPAE (ATCC CCL 209), Saos-2 (ATCC HTB-85), ARPE-19 human retinal pigment epithelium (ATCC CRL-2302), *Xenopus* melanophores, and *Xenopus* oocytes.

A variety of mammalian expression vectors can be used to express recombinant human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in mammalian cells. Commercially available mammalian expression vectors which are suitable include, but are not limited to, pMCIneo (Stratagene), pSG5 (Stratagene), pcDNAI and pcDNAIamp, pcDNA3, pcDNA3.1, pCR3.1 (Invitrogen), EBO-pSV2-neo (ATCC 37593), pBPV-1(8-2) (ATCC 37110), pdBPV-MMTneo(342-12) (ATCC 37224), pRSVgpt (ATCC 37199), pRSVneo (ATCC 37198), pIZD35 (ATCC 37565), and pSV2-dhfr (ATCC 37146). Another suitable vector is the PT7TS oocyte expression vector.

Following expression in recombinant cells, human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can be purified by conventional techniques to a level that is substantially free from other proteins. Techniques that can be used include ammonium sulfate precipitation, hydrophobic or hydrophilic interaction chromatography, ion exchange chromatography, affinity chromatography, phosphocellulose chromatography, size exclusion chromatography, preparative gel electrophoresis, and alcohol precipitation. In some cases, it may be

advantageous to employ protein denaturing and/or refolding steps in addition to such techniques.

- Certain voltage-gated potassium channel subunits have been found to require the expression of other voltage-gated potassium channel subunits in order to be properly expressed at high levels and inserted in membranes. For example, co-expression of KCNQ3 appears to enhance the expression of KCNQ2 in *Xenopus* oocytes (Wang et al., 1998, Science 282:1890-1893). Also, some voltage-gated potassium channel α subunits require other related α subunits (Jegla and Salkoff, 1997, J. Neurosci. 17:32-44) or Kv β 2 subunits (Shi et al., 1995, Neuron 16:843-852).
- Accordingly, the recombinant expression of the human calcium sensitive potassium channel β 2, β 3a, β 3b, β 3c, or β 3d subunit proteins may, under certain circumstances, benefit from the co-expression of other proteins and such co-expression is intended to be within the scope of the present invention. A particularly preferred form of co-expression is the co-expression of a human calcium sensitive potassium channel β 2, β 3a, β 3b, β 3c, or β 3d subunit protein (or combinations thereof) with a human calcium sensitive potassium channel α subunit protein. Such co-expression can be effected by transfecting an expression vector encoding a human calcium sensitive potassium channel β 2, β 3a, β 3b, β 3c, or β 3d subunit protein into a cell that naturally expresses a human calcium sensitive potassium channel α subunit protein.
- Alternatively, an expression vector encoding a human calcium sensitive potassium channel β 2, β 3a, β 3b, β 3c, or β 3d subunit protein can be transfected into a cell in which an expression vector encoding a human calcium sensitive potassium channel α subunit protein has also been transfected. Preferably, such a cell does not naturally express human calcium sensitive potassium channel α or β subunits.
- The present invention includes human calcium sensitive potassium channel β 2, β 3a, β 3b, β 3c, and β 3d subunit proteins substantially free from other proteins. The deduced amino acid sequences of the full-length human calcium sensitive potassium channel β 2, β 3a, β 3b, β 3c, and β 3d subunit proteins are shown in SEQ.ID.NO.:2, 4, 6, 8, and 10, respectively. Thus, the present invention includes human calcium sensitive potassium channel β 2, β 3a, β 3b, β 3c, and β 3d subunit proteins substantially free from other proteins having the amino acid sequences SEQ.ID.NO.:2, SEQ.ID.NO.:4; SEQ.ID.NO.:4 with an asparagine at position 163 instead of a serine; SEQ.ID.NO.:6; SEQ.ID.NO.:6 with a serine at position 143 instead of an asparagine; SEQ.ID.NO.:8; SEQ.ID.NO.:8 with an asparagine at

position 161 instead of a serine; SEQ.ID.NO.:10; and SEQ.ID.NO.:10 with a serine at position 165 instead of an asparagine. The present invention also includes isolated human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ subunit proteins having the amino acid sequences SEQ.ID.NO.:2, SEQ.ID.NO.:4;

- 5 SEQ.ID.NO.:4 with an asparagine at position 163 instead of a serine; SEQ.ID.NO.:6; SEQ.ID.NO.:6 with a serine at position 143 instead of an asparagine; SEQ.ID.NO.:8; SEQ.ID.NO.:8 with an asparagine at position 161 instead of a serine; SEQ.ID.NO.:10; and SEQ.ID.NO.:10 with a serine at position 165 instead of an asparagine.

- 10 Mutated forms of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ subunit proteins are intended to be within the scope of the present invention. In particular, mutated forms of SEQ.ID.NOs:2, 4, 6, 8, and 10 that give rise to calcium sensitive potassium channels having altered electrophysiological or pharmacological properties when combined with α subunits are within the scope of the present invention.

- 15 As with many proteins, it is possible to modify many of the amino acids of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins and still retain substantially the same biological activity as for the original proteins. Thus, the present invention includes modified human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ subunit proteins which have
20 amino acid deletions, additions, or substitutions but that still retain substantially the same biological activity as naturally occurring human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. It is generally accepted that single amino acid substitutions do not usually alter the biological activity of a protein (see, e.g., Molecular Biology of the Gene, Watson *et al.*, 1987, Fourth Ed., The
25 Benjamin/Cummings Publishing Co., Inc., page 226; and Cunningham & Wells, 1989, Science 244:1081-1085). Accordingly, the present invention includes polypeptides where one amino acid substitution has been made in SEQ.ID.NOs:2, 4, 6, 8, or 10 wherein the polypeptides still retain substantially the same biological activity as naturally occurring human calcium sensitive potassium channel $\beta 2$, $\beta 3a$,
30 $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. The present invention also includes polypeptides where two or more amino acid substitutions have been made in SEQ.ID.NOs:2, 4, 6, 8, or 10 wherein the polypeptides still retain substantially the same biological activity as naturally occurring human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. In particular, the present invention includes embodiments

where the above-described substitutions are conservative substitutions. In particular, the present invention includes embodiments where the above-described substitutions do not occur in conserved positions. Conserved positions are those positions in which the human calcium sensitive potassium channel $\beta 1$, $\beta 2$, and any of the $\beta 3$ subunits all
5 have the same amino acid (see Figure 6).

The human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins of the present invention may contain post-translational modifications, *e.g.*, covalently linked carbohydrate, phosphorylation, myristoylation, palmytoylation, *etc.*

10 The present invention also includes chimeric human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Chimeric human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins consist of a contiguous polypeptide sequence of at least a portion of a human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein fused to a
15 polypeptide sequence that is not from a human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein.

The present invention also includes isolated human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins and DNA encoding these isolated subunits. Use of the term "isolated" indicates that the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein or DNA has
20 been removed from its normal cellular environment. Thus, an isolated human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein may be in a cell-free solution or placed in a different cellular environment from that in which it occurs naturally. The term isolated does not imply that an isolated human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein is the
25 only protein present, but instead means that the isolated human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein is at least 95% free of non-amino acid material (*e.g.*, nucleic acids, lipids, carbohydrates) naturally associated with the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein. Thus, a human calcium sensitive potassium channel $\beta 2$, $\beta 3a$,
30 $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein that is expressed in bacteria or even in eukaryotic cells which do not naturally (*i.e.*, without human intervention) express it through recombinant means is an "isolated human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein."

It is known that certain potassium channels subunits can interact to form heteromeric structures resulting in functional potassium channels. For example, KCNQ2 and KCNQ3 can assemble to form a heteromeric functional potassium channel (Wang et al., 1998, Science 282:1890-1893). Accordingly, it is believed likely that the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins of the present invention will also be able to form heteromeric structures with other proteins where such heteromeric structures constitute functional potassium channels. Thus, the present invention includes such heteromers comprising human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Preferred heteromers are those in which the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins of the present invention forms heteromers with calcium sensitive potassium channel α subunits.

DNA encoding the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can be obtained by methods well known in the art. For example, a cDNA fragment encoding full-length human calcium sensitive potassium channel $\beta 2$ subunit protein can be isolated from human uterus, ovary or pancreas cDNA by using the polymerase chain reaction (PCR) employing suitable primer pairs. Such primer pairs can be selected based upon the DNA sequence encoding the human calcium sensitive potassium channel $\beta 2$ subunit protein shown in Figure 1A as SEQ.ID.NO.:1. Suitable primer pairs would be, *e.g.*:

5'-AAG ATG TTT ATA TGG ACC AGT GGC-3' (SEQ.ID.NO.:12)

and

5'-ACT CAT AAC AGA CTG CAC GTT AC-3' (SEQ.ID.NO.:13).

The above and subsequent primers are meant to be illustrative only; one skilled in the art would readily be able to design other suitable primers based upon SEQ.ID.NO.:1. Such primers could be produced by methods of oligonucleotide synthesis that are well known in the art.

In a similar manner, PCR primers can be selected and produced for the other human calcium sensitive potassium channel subunit proteins of the present invention. For example, for the human calcium sensitive potassium channel $\beta 3a$ subunit, suitable primer pairs would be, *e.g.*:

5'-GTC ATG CAG CCC TTC AGC ATC CC-3' (SEQ.ID.NO.:14)

and

5'-TTG CAG AAA TCA CAG ACA TCT GAA-3' (SEQ.ID.NO.:15).

A suitable cDNA template from which the human calcium sensitive potassium channel $\beta 3a$ subunit can be isolated is human heart, skeletal muscle or spleen cDNA.

5 For the human calcium sensitive potassium channel $\beta 3b$ subunit, suitable primer pairs would be, *e.g.*:

5'-GCA ATG ACA GCC TTT CCT GCC TC-3' (SEQ.ID.NO.:16)

and

5'-TTG CAG AAA TCA CAG ACA TCT GAA-3' (SEQ.ID.NO.:15).

10

A suitable cDNA template from which the human calcium sensitive potassium channel $\beta 3b$ subunit can be isolated is human spleen cDNA.

For the human calcium sensitive potassium channel $\beta 3c$ subunit, suitable primer pairs would be, *e.g.*:

15 5'-GAA ATG TTC CCC CTT CTT TAT GAG-3' (SEQ.ID.NO.:17)

and

5'-TTG CAG AAA TCA CAG ACA TCT GAA-3' (SEQ.ID.NO.:15).

20 A suitable cDNA template from which the human calcium sensitive potassium channel $\beta 3c$ subunit can be isolated is human pancreas or spleen cDNA.

For the human calcium sensitive potassium channel $\beta 3d$ subunit, suitable primer pairs would be, *e.g.*:

5'-GAG ATG GAC TTT TCA CCA AGC TCT-3' (SEQ.ID.NO.:18)

and

25 5'-TTG CAG AAA TCA CAG ACA TCT GAA-3' (SEQ.ID.NO.:15).

A suitable cDNA template from which the human calcium sensitive potassium channel $\beta 3d$ subunit can be isolated is human pancreas or spleen cDNA.

30 PCR reactions can be carried out with a variety of thermostable enzymes including but not limited to AmpliTaq, AmpliTaq Gold, or Vent polymerase. For AmpliTaq, reactions can be carried out in 10 mM Tris-Cl, pH 8.3, 2.0 mM $MgCl_2$, 200 μM for each dNTP, 50 mM KCl, 0.2 μM for each primer, 10 ng of DNA template, 0.05 units/ μl of AmpliTaq. The reactions are heated at 95°C for 3 minutes and then cycled 25 times using the cycling parameters of 95°C, 20 seconds, 62°C, 20

seconds, 72°C, 3 minutes. In addition to these conditions, a variety of suitable PCR protocols can be found in PCR Primer, A Laboratory Manual, edited by C.W. Dieffenbach and G.S. Dveksler, 1995, Cold Spring Harbor Laboratory Press; or PCR Protocols: A Guide to Methods and Applications, Michael *et al.*, eds., 1990,

5 Academic Press.

Since the calcium sensitive potassium channel subunits of the present invention are highly homologous to one another, and to other potassium channel subunits, it is desirable to sequence the clones obtained by the herein-described methods, in order to verify that the desired calcium sensitive potassium channel β subunits have in fact been obtained.

By these methods, cDNA clones encoding the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can be obtained. These cDNA clones can be cloned into suitable cloning vectors or expression vectors, *e.g.*, the mammalian expression vector pcDNA3.1 (Invitrogen, San Diego, CA). Human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can then be produced by transfecting expression vectors encoding the subunits or portions thereof into suitable host cells and growing the host cells under appropriate conditions. Human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can then be isolated by methods well known in the art.

As an alternative to the above-described PCR methods, cDNA clones encoding the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can be isolated from cDNA libraries using as a probe oligonucleotides specific for each human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit and methods well known in the art for screening cDNA libraries with oligonucleotide probes. Such methods are described in, *e.g.*, Sambrook *et al.*, 1989, *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory, Cold Spring Harbor, New York; Glover, D.M. (ed.), 1985, *DNA Cloning: A Practical Approach*, MRL Press, Ltd., Oxford, U.K., Vol. I, II. Oligonucleotides that are specific for particular human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits and that can be used to screen cDNA libraries can be readily designed based upon the DNA sequences shown in Figures 1-5 and can be synthesized by methods well-known in the art.

Genomic clones containing the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit genes can be obtained from commercially

available human PAC, YAC, or BAC libraries available from Research Genetics, Huntsville, AL. Alternatively, one may prepare genomic libraries, *e.g.*, in P1 artificial chromosome vectors, from which genomic clones containing the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit genes can be isolated, using probes based upon the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit DNA sequences disclosed herein. Methods of preparing such libraries are known in the art (see, *e.g.*, Ioannou *et al.*, 1994, Nature Genet. 6:84-89).

The novel DNA sequences of the present invention can be used in various diagnostic methods. The present invention provides diagnostic methods for determining whether a patient carries a mutation in one or more of the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit genes. In broad terms, such methods comprise determining the DNA sequence of a region in or near one or more of the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit genes from the patient and comparing that sequence to the sequence from the corresponding region of the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit genes from a non-affected person, *i.e.*, a person who does not have the condition which is being diagnosed, where a difference in sequence between the DNA sequence of the gene from the patient and the DNA sequence of the gene from the non-affected person indicates that the patient has a mutation in one or more of the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit genes.

The present invention also provides oligonucleotide probes, based upon the sequences of SEQ.ID.NOs: 1, 3, 5, 7, 9, or 20 that can be used in diagnostic methods to identify patients having mutated forms of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits, to determine the level of expression of RNA encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits, or to isolate genes homologous to human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits. In particular, the present invention includes DNA oligonucleotides comprising at least about 10, 15, or 18 contiguous nucleotides of a sequence selected from the group consisting of: SEQ.ID.NOs: 1, 3, 5, 7, 9, and 20 where the oligonucleotide probe comprises no stretch of contiguous nucleotides longer than 5 of a sequence selected from the group consisting of: SEQ.ID.NOs: 1, 3, 5, 7, 9, and 20 other than the said at least about 10,

15, or 18 contiguous nucleotides. The oligonucleotides can be substantially free from other nucleic acids. Also provided by the present invention are corresponding RNA oligonucleotides. The DNA or RNA oligonucleotide can be packaged in kits for use as probes.

5 The present invention makes possible the recombinant expression of the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in various cell types.

 The $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ subunits of the human calcium sensitive potassium channel have been expressed in *Xenopus* oocytes, both by themselves and in combination with an α subunit of a large-conductance calcium-sensitive potassium channel (maxi-K channel). The β subunits do not express currents on their own. However, when co-expressed with the α subunit, the $\beta 2$, $\beta 3a$, and $\beta 3c$ subunits induce inactivation of calcium sensitive potassium currents (Figure 7). The rates of inactivation produced by $\beta 2$, $\beta 3a$ and $\beta 3c$ are dependent upon voltage and internal calcium concentration; inactivation time constants reach a maximum at high depolarizations and high micromolar calcium for $\beta 2$, $\beta 3a$ and $\beta 3c$, $\tau_{inact} \sim 30-40$ ms at 80 mV with 30 μM intracellular Ca^{2+} . Measurements of current-voltage dependence obtained in the presence of micromolar intracellular Ca^{2+} demonstrate that $\beta 2$ subunits induce a large shift in the voltage dependence of activation (~ 80 mV towards negative potentials, with 30 μM Ca^{2+} in the bath; Figure 7B). This modulatory effect is similar to the one previously described for $\beta 1$ subunits, which do not induce inactivation. (McManus et al., 1995, Neuron 14:645-650). In contrast, $\beta 3a$, $\beta 3b$, $\beta 3c$, and $\beta 3d$ subunits do not shift the voltage dependence when compared to channels containing only α subunits (Figure 7B).

25 The present invention also makes possible the development of assays that measure the biological activity of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Such assays using recombinantly expressed human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are especially of interest. Such assays can be used to screen libraries of compounds or other sources of compounds to identify compounds that are activators or inhibitors of the activity of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Such identified compounds can serve as "leads" for the development of pharmaceuticals that can be used to treat

patients having diseases in which it is beneficial to enhance or suppress calcium sensitive potassium channel activity.

In versions of the above-described assays, calcium sensitive potassium channels containing mutant human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are used and inhibitors or activators of the activity of the mutant calcium sensitive potassium channels are identified.

Preferred cell lines for recombinant expression of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are those which do not express endogenous potassium channels (*e.g.*, CV-1, NIH-3T3, CHO-K1, COS-7). Such cell lines can be loaded with ^{86}Rb , an ion which can pass through potassium channels. The ^{86}Rb -loaded cells can be exposed to collections of substances (*e.g.*, combinatorial libraries, natural products, analogues of lead compounds produced by medicinal chemistry) and those substances that are able to alter ^{86}Rb efflux identified. Such substances are likely to be activators or inhibitors of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

Activators and inhibitors of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are likely to be substances that are capable of binding to calcium sensitive potassium channels. Accordingly, one type of assay determines whether one or more of a collection of substances is capable of such binding.

Accordingly, the present invention provides a method for identifying substances that bind to calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins comprising:

- (a) providing cells expressing a calcium sensitive potassium channel containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;
- (b) exposing the cells containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins to a substance that is not known to bind calcium sensitive potassium channels;
- (c) determining the amount of binding of the substance to the cells;
- (d) comparing the amount of binding in step (c) to the amount of binding of the substance to control cells where the control cells are substantially

identical to the cells of step (a) except that the control cells do not express human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

where if the amount of binding in step (c) is greater than the amount of binding of the substance to control cells, then the substance binds to calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

Another version of this assay makes use of compounds that are known to bind to calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. New binders are identified by virtue of their ability to potentiate, prevent, or displace the binding of the known compounds. Substances that have this ability are likely themselves to be inhibitors or activators of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

Accordingly, the present invention includes a method of identifying substances that bind calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins and thus are likely to be inhibitors or activators of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins comprising:

(a) providing cells expressing calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

(b) exposing the cells to a compound that is known to bind to the calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

(c) determining the amount of binding of the compound to the cells in the presence and in the absence of a substance not known to bind to calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

where if the amount of binding of the compound in the presence of the substance differs from that in the absence of the substance, then the substance binds calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins and is likely to be an inhibitor or

activator of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

Generally, the known compound is labeled (*e.g.*, radioactively, enzymatically, fluorescently) in order to facilitate measuring its binding to the

5 calcium sensitive potassium channels.

Once a substance has been identified by the above-described methods, it can be assayed in functional tests, such as those described herein, in order to determine whether it is an inhibitor or an activator.

In particular embodiments, the compound known to bind calcium
10 sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins is selected from the group consisting of: charybdotoxin, iberiotoxin, and dehydrosoyasaponin.

The present invention includes a method of identifying activators or inhibitors of calcium sensitive potassium channels containing human calcium
15 sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins comprising:

(a) recombinantly expressing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins or mutant human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in a host cell so that the recombinantly expressed human calcium sensitive potassium channel $\beta 2$,
20 $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins form calcium sensitive potassium channels by forming heteromers with other calcium sensitive potassium channel subunit proteins;

(b) measuring the biological activity of the calcium sensitive potassium channels formed in step (a) in the presence and in the absence of a substance suspected of being an activator or an inhibitor of calcium sensitive
25 potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

where a change in the biological activity of the calcium sensitive potassium channels formed in step (a) in the presence as compared to the absence of the substance indicates that the substance is an activator or an inhibitor of calcium
30 sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

It may be advantageous to recombinantly express other subunits of calcium sensitive potassium channels such as, *e.g.*, an α subunit. Alternatively, it may be advantageous to use host cells that endogenously express such other subunits.

In particular embodiments, the biological activity is the production of a calcium sensitive potassium current, a FRET signal, or the efflux of ^{86}Rb .

In particular embodiments, a vector encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins is transferred into
5 *Xenopus* oocytes in order to cause the expression of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in the oocytes. Alternatively, RNA encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can be prepared *in vitro* and injected into the oocytes, also resulting in the expression of human calcium sensitive potassium
10 channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in the oocytes. Following expression of the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in the oocytes, and following the formation of calcium sensitive potassium channels containing these subunits and other calcium sensitive potassium channel subunits (which other subunits may also be transferred into the oocytes),
15 membrane currents are measured after the transmembrane voltage and/or internal calcium concentration is changed in steps. A change in membrane current is observed when the calcium sensitive potassium channels open or close, allowing or inhibiting potassium ion flow, respectively. Similar oocyte studies were reported for KCNQ2 and KCNQ3 potassium channels in Wang et al., 1998, Science 282:1890-1893 and
20 this reference and references cited therein can be consulted for guidance as to how to carry out such studies.

Inhibitors or activators of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can be identified by exposing the oocytes to collections of substances
25 and determining whether the substances can block or diminish, or activate or enhance the membrane currents observed in the absence of the substance.

Accordingly, the present invention provides a method of identifying inhibitors or activators of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins
30 comprising:

(a) expressing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in a heterologous system such that calcium sensitive potassium channels containing the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are formed;

(b) changing the transmembrane potential or internal calcium concentration of the heterologous system in the presence and the absence of a substance suspected of being an inhibitor or activator of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

(c) measuring membrane potassium currents following step (b);
where if the membrane potassium currents measured in step (c) are greater in the absence rather than in the presence of the substance, then the substance is an inhibitor of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

where if the membrane potassium currents measured in step (c) are less in the absence rather than in the presence of the substance, then the substance is an activator of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

In particular embodiments, the heterologous system is selected from the group consisting of: *Xenopus* oocytes and a mammalian cell line.

The present invention also includes assays for the identification of activators and inhibitors of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins that are based upon fluorescence resonance energy transfer (FRET) between a first and a second fluorescent dye where the first dye is bound to one side of the plasma membrane of a cell expressing calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins and the second dye is free to shuttle from one face of the membrane to the other face in response to changes in membrane potential. In certain embodiments, the first dye is impenetrable to the plasma membrane of the cells and is bound predominately to the extracellular surface of the plasma membrane. The second dye is trapped within the plasma membrane but is free to diffuse within the membrane. At normal (*i.e.*, negative) resting potentials of the membrane, the second dye is bound predominately to the inner surface of the extracellular face of the plasma membrane, thus placing the second dye in close proximity to the first dye. This close proximity allows for the generation of a large amount of FRET between the two dyes. Following membrane depolarization, the second dye moves from the extracellular face of the membrane to the intracellular face, thus increasing the distance between

the dyes. This increased distance results in a decrease in FRET, with a corresponding increase in fluorescent emission derived from the first dye and a corresponding decrease in the fluorescent emission from the second dye. In this way, the amount of FRET between the two dyes can be used to measure the polarization state of the membrane. For a fuller description of this technique, see González & Tsien, 1997, Chemistry & Biology 4:269-277. See also González & Tsien, 1995, Biophys. J. 69:1272-1280 and U.S. Patent No. 5,661,035.

In certain embodiments, the first dye is a fluorescent lectin or a fluorescent phospholipid that acts as the fluorescent donor. Examples of such a first dye are: a coumarin-labeled phosphatidylethanolamine (*e.g.*, N-(6-chloro-7-hydroxy-2-oxo-2H--1-benzopyran-3-carboxamidoacetyl)-dimyristoylphosphatidylethanolamine) or N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)-dipalmitoylphosphatidylethanolamine); a fluorescently-labeled lectin (*e.g.*, fluorescein-labeled wheat germ agglutinin). In certain embodiments, the second dye is an oxonol that acts as the fluorescent acceptor. Examples of such a second dye are: bis(1,3-dialkyl-2-thiobarbiturate)trimethineoxonols (*e.g.*, bis(1,3-dihexyl-2-thiobarbiturate)trimethineoxonol) or pentamethineoxonol analogues (*e.g.*, bis(1,3-dihexyl-2-thiobarbiturate)pentamethineoxonol; or bis(1,3-dibutyl-2-thiobarbiturate)pentamethineoxonol). See González & Tsien, 1997, Chemistry & Biology 4:269-277 for methods of synthesizing various dyes suitable for use in the present invention. In certain embodiments, the assay may comprise a natural carotenoid, *e.g.*, astaxanthin, in order to reduce photodynamic damage due to singlet oxygen.

The above described assays can be utilized to discover activators and inhibitors of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Such assays will generally utilize cells that express calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins, *e.g.*, by transfection with expression vectors encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins and, optionally, other calcium sensitive potassium channel subunits. In such cells, depolarization of the membrane potential as well as increases in intracellular calcium concentration will tend to open the calcium sensitive potassium channels. This will result in potassium efflux, tending to counteract the depolarization. In other words,

the cells will tend to repolarize. The presence of an inhibitor of the calcium sensitive potassium channel will prevent, or diminish, this repolarization. Thus, membrane potential will tend to become more positive (*i.e.*, depolarized) in the presence of inhibitors. Activators of the calcium sensitive potassium channel will open this
5 channel and thus tend to hyperpolarize the membrane potential. Changes in membrane potential (depolarizations and hyperpolarizations) that are caused by inhibitors and activators of calcium sensitive potassium channels can be monitored by the assays using FRET described above.

Accordingly, the present invention provides a method of identifying
10 activators of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins comprising:

- (a) providing test cells comprising:
 - (1) an expression vector that directs the expression of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit
15 proteins in the cells so that calcium sensitive potassium channels containing human $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are formed in the cells;
 - (2) a first fluorescent dye, where the first dye is bound to one side of the plasma membrane of the cells; and
 - (3) a second fluorescent dye, where the second fluorescent
20 dye is free to shuttle from one face of the plasma membrane of the cells to the other face in response to changes in membrane potential;
- (b) exposing the test cells to a substance that is suspected of being an activator of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;
- 25 (c) measuring the amount of fluorescence resonance energy transfer (FRET) in the test cells that have been exposed to the substance;
- (d) comparing the amount of FRET exhibited by the test cells that have been exposed to the substance with the amount of FRET exhibited by control cells;
- 30 wherein if the amount of FRET exhibited by the test cells is greater than the amount of FRET exhibited by the control cells, the substance is an activator of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ -subunit proteins;

where the control cells are either (1) cells that are essentially the same as the test cells except that they do not comprise at least one of the items listed at (a) (1)-(3) but have been exposed to the substance; or (2) test cells that have not been exposed to the substance.

5 The present invention also provides a method of identifying inhibitors of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins comprising:

- (a) providing test cells comprising:
 - (1) an expression vector that directs the expression of
 - 10 human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in the cells so that calcium sensitive potassium channels containing human $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are formed in the cells;
 - (2) a first fluorescent dye, where the first dye is bound to one side of the plasma membrane of the cells; and
 - 15 (3) a second fluorescent dye, where the second fluorescent dye is free to shuttle from one face of the plasma membrane of the cells to the other face in response to changes in membrane potential;
- (b) exposing the test cells to a substance that is suspected of being an inhibitor of calcium sensitive potassium channels containing human calcium
- 20 sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;
- (c) measuring the amount of fluorescence resonance energy transfer (FRET) in the test cells that have been exposed to the substance;
- (d) comparing the amount of FRET exhibited by the test cells that have been exposed to the substance with the amount of FRET exhibited by control
- 25 cells;

wherein if the amount of FRET exhibited by the test cells is less than the amount of FRET exhibited by the control cells, the substance is an inhibitor of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

30 where the control cells are either (1) cells that are essentially the same as the test cells except that they do not comprise at least one of the items listed at (a) (1)-(3) but have been exposed to the substance; or (2) test cells that have not been exposed to the substance.

In a variation of the assay described above, instead of the cell's membrane potential being allowed to reach steady state on its own, the membrane potential is artificially set at a potential in which the calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are open. This can be done, *e.g.*, by variation of the external K^+ concentration in a known manner (*e.g.*, increased concentrations of external K^+). If such cells, having open calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins, are exposed to inhibitors, the calcium sensitive potassium channels will be blocked, and the cells' membrane potentials will be depolarized. This depolarization can be observed as a decrease in FRET.

In a variation of the assay described above, instead of the cell's membrane potential being allowed to reach steady state on its own, the membrane potential is artificially set at a potential in which the calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are open by coexpression of another depolarizing current. If such cells, having open calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins, are exposed to inhibitors, the calcium sensitive potassium channels will be blocked, and the cells' membrane potentials will be depolarized. This depolarization can be observed as a decrease in FRET. If such cells, having open calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins, are exposed to activators, the balance of the calcium sensitive potassium current and the additional depolarizing current will shift (*i.e.*, the calcium sensitive potassium current will make a larger contribution to the total current) and the cell's membrane potential will be hyperpolarized. This polarization may be observed as an increase in FRET.

Accordingly, the present invention provides a method of identifying inhibitors or activators of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins comprising:

- (a) providing cells comprising:
 - (1) an expression vector that directs the expression of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit

proteins in the cells so that calcium sensitive potassium channels containing human $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins are formed in the cells;

(2) a first fluorescent dye, where the first dye is bound to one side of the plasma membrane of the cells; and

5 (3) a second fluorescent dye, where the second fluorescent dye is free to shuttle from one face of the plasma membrane of the cells to the other face in response to changes in membrane potential;

(b) adjusting the membrane potential of the cells such that the ion channel formed by the calcium sensitive potassium channels containing human
10 calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins is open;

(c) measuring the amount of fluorescence resonance energy transfer (FRET) in the test cells;

(d) repeating step (b) and step (c) while the cells are exposed to a
15 substance that is suspected of being an inhibitor or activator of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

where if the amount of FRET exhibited by the cells that are exposed to the substance is different than the amount of FRET exhibited by the cells that have
20 not been exposed to the substance, then the substance is an inhibitor or activator of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

In particular embodiments of the above-described methods, the cells contain an expression vector encoding a human calcium sensitive potassium channel
25 $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein. In particular embodiments of the above-described methods, the expression vector is transfected into the test cells.

In particular embodiments of the above-described methods, the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein has an amino acid sequence selected from the group consisting of: SEQ.ID.NO.:2, 4, 6, 8,
30 and 10.

In particular embodiments of the above-described methods, the first fluorescent dye is selected from the group consisting of: a fluorescent lectin; a fluorescent phospholipid; a coumarin-labeled phosphatidylethanolamine; N-(6-chloro-7-hydroxy-2-oxo-2H--1-benzopyran-3-carboxamidoacetyl)-dimyristoylphosphatidyl-

ethanolamine); N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)-dipalmitoyl-phosphatidylethanolamine); and fluorescein-labeled wheat germ agglutinin.

In particular embodiments of the above-described methods, the second fluorescent dye is selected from the group consisting of: an oxonol that acts as the
 5 fluorescent acceptor; bis(1,3-dialkyl-2-thiobarbiturate)trimethineoxonols; bis(1,3-dihexyl-2-thiobarbiturate)trimethineoxonol; bis(1,3-dialkyl-2-thiobarbiturate)quatramethineoxonols; bis(1,3-dialkyl-2-thiobarbiturate)pentamethineoxonols; bis(1,3-dihexyl-2-thiobarbiturate)pentamethineoxonol; bis(1,3-dibutyl-2-thiobarbiturate)pentamethineoxonol); and bis(1,3-dialkyl-2-thiobarbiturate)
 10 hexamethineoxonols.

In a particular embodiment of the above-described methods, the cells are eukaryotic cells. In another embodiment, the cells are mammalian cells. In other embodiments, the cells are L cells L-M(TK⁻) (ATCC CCL 1.3), L cells L-M (ATCC CCL 1.2), 293 (ATCC CRL 1573), Raji (ATCC CCL 86), CV-1 (ATCC CCL 70),
 15 COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCC CCL 2), C1271 (ATCC CRL 1616), BS-C-1 (ATCC CCL 26), MRC-5 (ATCC CCL 171), *Xenopus* melanophores, or *Xenopus* oocytes.

In particular embodiments of the above-described methods, the control
 20 cells do not comprise item (a)(1) but do comprise items (a)(2) and (a)(3).

In assays to identify activators or inhibitors of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins, it may be advantageous to co-express another calcium sensitive potassium channel subunit besides the human calcium sensitive
 25 potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit. In particular, it may be advantageous to co-express a calcium sensitive potassium channel α subunit. Preferably, this is done by co-transfecting into the cells an expression vector encoding the other subunit. Suitable other subunits are, *e.g.*, the human calcium sensitive potassium channel α subunit *h-slo* (GenBank accession no. U11058), the mouse
 30 calcium sensitive potassium channel α subunit *m-slo* (GenBank accession no. U09383), the small conductance calcium sensitive potassium α subunits (GenBank accession nos. U69883, U69882, AF031815), or the intermediate conductance calcium sensitive potassium channel α subunit (GenBank accession no. AF022797).

Small regions of genomic sequences in proximity to a gene often regulate the transcription of that gene. These sequences are referred to as cis-acting elements. The proteins that bind these DNA sequences and directly affect the ability of the transcriptional machinery to bind or transcribe the gene are referred to as trans-acting elements. The cis-acting transcriptional regulatory elements are most often 5' of the transcription start site, but have been located within and 3' of the transcribed portion of genes as well. Depending on their effects on the rate of transcription, these sequences can be divided into three categories: promoters, enhancers, and repressors. A promoter independently allows transcription of the gene, while an enhancer increases the rate of transcription but is not capable of inducing transcription independently of the promoter. A repressor element inhibits transcription directed by a promoter element. Methods for identifying these elements are well known in the field and are described in Ausubel et al., eds., 1989, Current Protocols in Molecular Biology, sections 9.6-9.8, and 12.0-12.11, John Wiley & Sons, New York, NY.

Accordingly, the novel genomic sequences (SEQ.ID.NO.:20, Figure 8) and isolated BAC clones of the present invention make possible methods for identifying 1) DNA sequences required for transcriptional control of gene expression, 2) proteins involved in transcriptional regulation and 3) compounds which modulate the rate of transcription of the $\beta 3$ gene. Such assays utilize isolated and/or recombinant DNA comprising portions of SEQ.ID.NO.:20, positions 1 to 17,436, inserted into vectors upstream of the open reading frame of a reporter protein.

Useful reporter proteins are ones that are not expressed in the cells to be assayed (or are easily distinguished from endogenous proteins), have a linear relationship between the abundance of the transcript and the abundance of the reporter protein, and have a large window between the minimum detection level and saturation of detection system. Ideally, the abundance of the reporter protein is quickly measured by an enzymatic reaction, fluorescence detection, immunoassay or other means. Typical reporter proteins include, but are not limited to, the following: Chloramphenicol Acetyltransferase (CAT), firefly luciferase, Beta-Lactamase, Beta-Galactosidase, Secreted Alkaline Phosphatase (SEAP), human Growth Hormone (hGH), Green Fluorescent Protein (GFP) and GFP derivatives. Reporter vectors incorporating these proteins are commercially available, as are similar reporter vectors containing constitutive promoters, enhancers, or both (Clontech).

The present invention provides a method for identifying nucleotide sequences involved in transcriptional regulation of $\beta 3$ gene expression. Once a fragment of at least 6 contiguous nucleotides of DNA from SEQ.ID.NO.:20, positions 1-17,436, has been inserted upstream of the reporter cDNA in a promoter-reporter vector, the vector is then transfected into cells that either do or do not endogenously express one or more of the calcium sensitive potassium channel subunits $\beta 3a$, $\beta 3b$, $\beta 3c$ or $\beta 3d$. Promoter-reporter vectors may contain promoters, enhancers, both, or neither. Transfected cells are then assayed for the amount of reporter protein present. Because both transfection efficiency and transcription rate directly affect reporter protein level, it is useful in these assays to determine the transfection efficiency by co-transfecting a second vector (molar ratio 1:1) containing a distinct reporter behind a constitutive promoter, and determining the fraction of transfected cells.

In versions of the above assay, vectors are constructed with fragments of SEQ.ID.NO.:20 inserted upstream of a reporter cDNA with no other enhancer or promoter elements. These vectors (with and without fragments of SEQ.ID.NO.:20) are transfected into cells that endogenously express $\beta 3$ subunits. Calcium sensitive potassium channel $\beta 3$ subunit promoter elements are identified by the ability of these 5' gene fragments to stimulate reporter expression above the levels observed in the parent vector. The minimum required promoter sequence is then identified by successively deleting regions of the identified promoter fragment, and repeating the assay.

Another version of the assay incorporates fragments of SEQ.ID.NO.:20 inserted upstream of the reporter cDNA in a promoter-reporter vector containing an enhancer element. These vectors (with and without fragments of SEQ.ID.NO.:20) are transfected into cells that endogenously express $\beta 3$ subunits. Weak calcium sensitive potassium channel $\beta 3$ subunit promoter elements are identified by the ability of these 5' gene fragments to stimulate reporter expression above the levels observed in the parent vector. The minimum required weak promoter sequence can then be identified by successively deleting regions of the identified weak promoter fragment and repeating the assay.

A different version of the assay incorporates fragments of SEQ.ID.NO.:20 inserted upstream of the reporter cDNA in a promoter-reporter vector with a constitutive promoter upstream. These vectors (with and without fragments of SEQ.ID.NO.:20) are transfected into cells that do not endogenously express $\beta 3$

subunits. Calcium sensitive potassium channel $\beta 3$ subunit repressor elements are identified by the ability of these 5' gene fragments to prevent or reduce reporter expression below the levels observed in the parent vector. The minimum required repressor sequence is then identified by successively deleting regions of the identified
5 repressor fragment and repeating the assay.

In view of the above, the present invention provides a method of identifying DNA sequences in the $\beta 3$ gene that promote, enhance, or repress gene transcription comprising:

- (a) constructing a promoter-reporter vector such that fragments of
10 the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) precede the coding cDNA sequence of a reporter gene which encodes a reporter protein;
- (b) transfecting the vector into cells and measuring the abundance of the reporter protein encoded by the vector;
- (c) comparing the abundance of the reporter protein in the cells of
15 step (b) to the abundance of the reporter protein in cells transfected with the vector without fragments of the promoter region of the $\beta 3$ gene;

where fragments of the promoter region of the $\beta 3$ gene which increase the abundance of the reporter protein in the absence of other promoter elements only in cells which endogenously express $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits are promoter
20 elements; sequences which decrease the abundance of the reporter protein in the presence of an unrelated constitutive promoter element in cells which do not endogenously express $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits are repressor elements; and sequences which increase the abundance of the reporter protein in the presence of an unrelated constitutive promoter element in cells which endogenously express $\beta 3a$,
25 $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits are enhancer elements.

In particular embodiments, the vector contains promoter or enhancer sequence elements which function independently of the fragments of the promoter region of the $\beta 3$ gene.

In particular embodiments, the abundance of the reporter protein is
30 normalized with respect to the fraction of transfected cells.

The binding of nuclear proteins to these sequences can be confirmed by gel-shift assays. A radiolabeled DNA fragment corresponding to the minimal sequence required to affect transcription is incubated with nuclear protein extracts from cells used to identify the regulatory DNA element, or tissues endogenously

expressing $\beta 3$ subunits. If a protein factor binds that sequence, the mobility in a gel will be altered, resulting in an apparent shift in the size of the radiolabeled fragment.

Transcription factors often are able to recognize more than one specific nucleotide sequence. As such, variations of sequences identified as minimal
5 promoters, enhancers or repressors necessary for transcriptional regulation of the $\beta 3$ gene in SEQ.ID.NO.:20, positions 1-17,436, which retain the ability to influence transcription as detected in the above described assays are intended to be included in the present invention.

Minimal promoter, enhancer or repressor DNA fragments thus
10 identified can then be used to identify and/or isolate proteins that influence transcriptional activity of the $\beta 3$ gene. Several methods are well known in the field, some of which are described in Ausubel et al., eds., 1989, Current Protocols in Molecular Biology, sections 12.0-12.11, John Wiley & Sons, New York, NY.

In one method, gel shift assays described above can be performed with
15 cloned or purified known transcription factors, to determine if they are capable of binding sequences involved in transcriptional regulation. Alternatively, super-shift assays can be performed in which an antibody that recognizes a particular transcription factor is added to the transcription factor-DNA complex. If the antibody binds to the transcription factor, which in turn binds the radiolabeled DNA fragment,
20 the mobility of the complex in a gel is further altered, resulting in a super-shift compared to the DNA alone. Using antibodies that recognize a specific transcription factor, or a class of transcription factors, allows identification of the factors involved in $\beta 3$ gene regulation. Variations of sequences identified as minimal promoters, enhancers or repressors necessary for transcriptional regulation of the $\beta 3$ gene in
25 SEQ.ID.NO.:20, positions 1-17,436, which retain the ability to undergo gel shifts or super-shifts as described in the above assays are intended to be included in the present invention.

In view of the above, the present invention provides a method of identifying DNA sequences in the $\beta 3$ gene that promote, enhance, or repress gene
30 transcription comprising:

(a) incubating radiolabeled fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) with nuclear extracts from cells and separating the incubation on a gel;

where fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene that migrate differently in a gel ('undergo a shift') after incubation with nuclear extracts from cells are DNA sequences which bind nuclear factors which promote, enhance or repress $\beta 3$ gene expression.

5 In particular embodiments, the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the method of claim 18.

 In particular embodiments, the cells express $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits.

10 In particular embodiments, the cells do not express $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits.

 The present invention provides a method of identifying nuclear factors involved in $\beta 3$ gene transcription regulation comprising:

 (a) incubating radiolabeled fragments of double stranded DNA
15 corresponding to sequences found in the promoter region of the $\beta 3$ gene
(SEQ.ID.NO.:20, nucleotides 1 to 17,436) with cloned or purified transcription factors and separating the incubation on a gel;

 where factors which bind $\beta 3$ gene promoter sequence elements will induce a shift in the migration of the radiolabeled DNA fragments, and are involved
20 in $\beta 3$ gene transcription regulation.

 In particular embodiments, the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

 The present invention provides a method of identifying nuclear factors
25 involved in $\beta 3$ gene transcription regulation comprising:

 (a) incubating radiolabeled fragments of double stranded DNA
corresponding to sequences found in the promoter region of the $\beta 3$ gene
(SEQ.ID.NO.:20, nucleotides 1 to 17,436) with nuclear extracts from cells and
separating the incubation on a gel;

30 (b) adding an antibody that specifically recognizes a single transcription factor or a family of transcription factors to the incubation of step (a), followed by separating the incubation on a gel;

where a super-shift in mobility of the double stranded DNA in step (b) as compared to step (a) indicates that a transcription factor recognized by the antibody binds the double stranded DNA.

In another method, the transcription factors that bind SEQ.ID.NO.:20,
5 positions 1-17,436, and regulate transcription can be purified. DNA fragments corresponding to the minimal sequence required to affect transcription are covalently linked to a matrix (typically an agarose bead). This matrix is then incubated with nuclear extracts of cells that contain factors which bind the minimal element. The matrix is then washed free of non-specific proteins and the factor(s) are eluted with an
10 excess of the DNA element, or by denaturation. Purified proteins can then be identified by immunoassay, protein sequencing, or other means.

Accordingly, the present invention provides a method of identifying nuclear factors involved in $\beta 3$ gene transcription regulation comprising:

- (a) attaching fragments of double stranded DNA corresponding to
15 sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) to a stable matrix;
- (b) incubating nuclear extracts from cells with the matrix;
- (c) washing non-binding proteins from the nuclear extract from the matrix;
- 20 (d) eluting bound proteins from the matrix with excess double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene;

where the eluted proteins from step (d) are nuclear factors involved in $\beta 3$ gene transcription regulation.

25 In particular embodiments, the method further comprises separating the eluted proteins from step (d) on a gel and staining the gel to test for purity of the eluted proteins.

In particular embodiments, the method further comprises sequencing the proteins that have been separated on the gel.

30 In particular embodiments, the method further comprises immunological analysis of the proteins that have been separated on the gel with antibodies directed towards known transcription factors to identify the eluted proteins by western blot or immunoprecipitation.

In particular embodiments, the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

5 In a different approach, cDNAs encoding the transcription factors that bind SEQ.ID.NO.:20; positions 1-17,436 can be cloned by several methods. In one version, the minimal DNA sequence is radiolabeled and used to screen an expression library made from tissues or cell lines that endogenously express the $\beta 3$ gene. Phage containing cDNA encoding the transcription factor are induced to express fusion proteins that target the transcription factor to its surface. Such phage plaques are
10 identified by their ability to bind radiolabeled DNA sequences containing the minimal DNA sequence.

Accordingly, the present invention provides a method of identifying clones encoding nuclear factors involved in $\beta 3$ gene transcription regulation by cloning comprising:

- 15 (a) screening an expression library with radiolabeled fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436)
- (b) determining which clones of the library bind the radiolabeled fragments of double stranded DNA;
- 20 (c) amplifying and sequencing the clones of step (b).

In particular embodiments, the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

Another cloning approach involves phage expressing transcription
25 factor fusion proteins at their surface. In this approach, the minimal DNA sequence is linked to a matrix. A phage expression library is then passed over the matrix and washed. Only phage containing the transcription factor bind the matrix. Bound phage are eluted with excess minimal DNA sequence and purified. cDNA encoding the transcription factor is then isolated from the phage and sequenced.

30 Accordingly, the present invention provides a method of identifying nuclear factors involved in $\beta 3$ gene transcription regulation by cloning comprising:

- (a) attaching fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) to a stable matrix;

(b) incubating phage expressing cDNA encoded fusion proteins at their surface with the matrix;

(c) removing phage that do not bind to the matrix by washing;

(d) eluting phage bound to the matrix with excess fragments of
5 double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene;

where the phage eluted in step (d) encode nuclear factors involved in $\beta 3$ gene transcription regulation.

In particular embodiments, the DNA corresponding to sequences found
10 in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

In particular embodiments, the phage eluted at step (d) are amplified and sequenced.

A separate transcription factor cloning approach is the yeast 'one-hybrid' method (available in kit form from Clontech). In this method, yeast strains
15 are made that contain several copies (three suggested) of the minimal element upstream of a reporter. A cDNA library is made such that each vector contains a cDNA that will be expressed as a fusion protein with the transcription activation domain of a yeast promoter. Thus, any fusion protein that specifically binds the DNA of interest will induce expression of the reporter protein. The vector containing the
20 cDNA is then isolated from the yeast and sequenced.

Accordingly, the present invention provides a method of identifying nuclear factors involved in $\beta 3$ gene transcription regulation by cloning comprising:

(a) constructing a yeast strain that contains a few to several copies of a fragment of double stranded DNA corresponding to sequences found in the
25 promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) preceding a cDNA encoding a reporter protein;

(b) constructing a cDNA library from cells in a vector that allows formation of fusion proteins encoded by the inserted cDNA and a transcription activation domain;

(c) transforming the library of (b) into the yeast strain of (a) and
30 isolating colonies of yeast displaying expression of the reporter protein.

In particular embodiments, the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

In particular embodiments, the method further comprises purifying the vectors from the isolated colonies and sequencing the cDNA in the vectors.

Since transcription factors often are able to recognize more than one specific nucleotide sequence, variations of sequences identified as minimal
5 promoters, enhancers or repressors necessary for transcriptional regulation of the $\beta 3$ gene in SEQ.ID.NO.:20; positions 1-17,436, that can be bound by transcription factors as detected in the above described assays are intended to be included in the present invention.

Identification of nucleotide sequences involved in transcriptional
10 regulation of $\beta 3$ gene expression by the methods described above allows for the development of assays that can be used to screen collections of substances to identify those substances that enhance or inhibit transcription of the $\beta 3$ gene. Fragments of the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) that have been shown to be involved in transcriptional regulation are linked to the coding
15 sequence of a reporter gene in a suitable vector and are then transferred to appropriate cells. The abundance of the reporter protein in the cells is determined. The cells are then exposed to compounds that are suspected of being capable of enhancing or inhibiting the rate of transcription of the $\beta 3$ gene. If the compound actually is capable of enhancing the rate of transcription of the $\beta 3$ gene, then the abundance of the
20 reporter protein will be increased when the cells are exposed to the compound. Conversely, if the compound actually is capable of inhibiting the rate of transcription of the $\beta 3$ gene, then the abundance of the reporter protein will be decreased when the cells are exposed to the compound.

Accordingly, the present invention provides a method of identifying
25 substances that enhance or inhibit the rate of transcription of the $\beta 3$ gene comprising:
(a) constructing a promoter-reporter vector such that fragments of the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) precede the coding cDNA sequence of a reporter gene which encodes a reporter protein;
(b) transfecting the vector into cells and measuring the abundance
30 of the reporter protein encoded by the vector in the presence and absence of a compound;

where (1) if the presence of the compound decreases the abundance of the reporter protein, then the compound is a substance that inhibits the rate of transcription of the $\beta 3$ gene; (2) if the presence of the compound increases the

abundance of the reporter protein, then the compound is a substance that enhances the rate of transcription of the $\beta 3$ gene.

In particular embodiments, the method further comprises a control in which the effect of the compound on the abundance of the reporter protein in control
5 cells is measured, where the control cells are cells that are essentially the same as the cells of step (b) except that the control cells have been transfected with a vector that lacks fragments of the promoter region of the $\beta 3$ gene.

While the above-described methods are explicitly directed to testing whether “a” substance is an activator or inhibitor of the transcription the $\beta 3$ gene or
10 the function of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins, it will be clear to one skilled in the art that such methods can be adapted to test collections of substances, *e.g.*, combinatorial libraries, to determine whether any members of such collections are activators or inhibitors of calcium sensitive potassium channels
15 containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Accordingly, the use of collections of substances, or individual members of such collections, as the substance in the above-described methods is within the scope of the present invention. In particular, it is envisioned that libraries that have been designed to incorporate chemical structures that are known to be
20 associated with potassium ion channel modulation, *e.g.*, dihydrobenzopyran libraries for potassium channel activators (International Patent Publication WO 95/30642) or biphenyl-derivative libraries for potassium channel inhibitors (International Patent Publication WO 95/04277) will be especially suitable.

The present invention includes pharmaceutical compositions
25 comprising activators or inhibitors of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins that have been identified by the herein-described methods as well as activators or inhibitors of $\beta 3$ gene transcription. The activators or inhibitors are generally combined with pharmaceutically acceptable carriers to form pharmaceutical compositions. Examples of such carriers and
30 methods of formulation of pharmaceutical compositions containing activators or inhibitors and carriers can be found in Remington’s Pharmaceutical Sciences. To form a pharmaceutically acceptable composition suitable for effective administration, such compositions will contain a therapeutically effective amount of the activators or inhibitors.

Therapeutic or prophylactic compositions are administered to an individual in amounts sufficient to treat or prevent conditions where human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein activity is abnormal. The effective amount can vary according to a variety of factors such as the individual's condition, weight, gender, and age. Other factors include the mode of administration. The appropriate amount can be determined by a skilled physician. Generally, an effective amount will be from about 0.01 to about 1,000, preferably from about 0.1 to about 250 and even more preferably from about 1 to about 50 mg per adult human per day.

Compositions can be used alone at appropriate dosages. Alternatively, co-administration or sequential administration of other agents can be desirable.

The compositions can be administered in a wide variety of therapeutic dosage forms in conventional vehicles for administration. For example, the compositions can be administered in such oral dosage forms as tablets, capsules (each including timed release and sustained release formulations), pills, powders, granules, elixirs, tinctures, solutions, suspensions, syrups and emulsions, or by injection. Likewise, they can also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts.

Compositions can be administered in a single daily dose, or the total daily dosage can be administered in divided doses of two, three, four or more times daily. Furthermore, compositions can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

The dosage regimen utilizing the compositions is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal, hepatic and cardiovascular function of the patient; and the particular composition thereof employed. A physician of ordinary skill can readily determine and prescribe the effective amount of the composition required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentrations of composition within the range that yields efficacy without

toxicity requires a regimen based on the kinetics of the composition's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a composition.

The inhibitors and activators of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins, or inhibitors and activators of $\beta 3$ subunit transcription will be useful for treating a variety of diseases involving excessive or insufficient calcium sensitive potassium channel activity. Accordingly, the present invention includes a method of treating asthma, diabetes, glaucoma, pregnant human myometrium, cerebral ischemia, and conditions where stimulation of neurotransmitter release is desired such as Alzheimer's disease and stimulation of damaged nerves by administering to a patient a therapeutically effective amount of a substance that is an activator or an inhibitor of a calcium sensitive potassium channel containing a human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein, or an activator or an inhibitor of $\beta 3$ subunit transcription.

The modulators of channel function or transcription activity of the present invention are also expected to be useful in conditions where currently marketed inhibitors of potassium channels such as glyburide, glipizide, and tolbutamide are useful, *e.g.*, as antidiabetic agents. Calcium sensitive potassium channels contribute to the repolarization, and thus the de-excitation, of neurons. Thus, inhibitors of calcium sensitive potassium channels are expected to act as agents that tend to keep neurons in a depolarized, excited state. Many diseases, such as depression and memory disorders are thought to result from the impairment of neurotransmitter release. As agents that contribute to neuronal excitability, the inhibitors of the present invention are expected to be useful in the treatment of such diseases since they will contribute to neuronal excitation and thus stimulate the release of neurotransmitters.

The activators of the present invention should be useful in conditions where it is desirable to decrease neuronal activity. Such conditions include, *e.g.*, excessive smooth muscle tone, angina, asthma, hypertension, incontinence, pre-term labor, migraine, cerebral ischemia, and Irritable Bowel Syndrome.

The calcium sensitive potassium channel subunits of the present invention are useful in conjunction with screens designed to identify activators and inhibitors of other ion channels. When screening compounds in order to identify

potential pharmaceuticals that specifically interact with a target ion channel, it is necessary to ensure that the compounds identified are as specific as possible for the target ion channel. To do this, it is necessary to screen the compounds against as wide an array as possible of ion channels that are similar to the target ion channel.

5 Thus, in order to find compounds that are potential pharmaceuticals that interact with ion channel A, it is not enough to ensure that the compounds interact with ion channel A (the “plus target”) and produce the desired pharmacological effect through ion channel A. It is also necessary to determine that the compounds do not interact with ion channels B, C, D, *etc.* (the “minus targets”). In general, as part of a screening
10 program, it is important to have as many minus targets as possible (see Hodgson, 1992, *Bio/Technology* 10:973-980, at 980). Human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins, DNA encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins, and recombinant cells that have been engineered to express human calcium sensitive
15 potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins have utility in that they can be used as “minus targets” in screens designed to identify compounds that specifically interact with other ion channels. For example, Wang et al., 1998, *Science* 282:1890-1893 have shown that KCNQ2 and KCNQ3 form a heteromeric potassium ion channel known as the “M-channel.” The M-channel is an important target for drug
20 discovery since mutations in KCNQ2 and KCNQ3 are responsible for causing epilepsy (Biervert et al., 1998, *Science* 279:403-406; Singh et al., 1998, *Nature Genet.* 18:25-29; Schroeder et al., *Nature* 1998, 396:687-690). A screening program designed to identify activators or inhibitors of the M-channel would benefit greatly by the use of potassium channels comprising human calcium sensitive potassium channel
25 $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins as minus targets.

The present invention also includes antibodies to the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins. Such antibodies may be polyclonal antibodies or monoclonal antibodies. The antibodies of the present invention can be raised against the entire human calcium sensitive
30 potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins or against suitable antigenic fragments of the subunit proteins that are coupled to suitable carriers, *e.g.*, serum albumin or keyhole limpet hemocyanin, by methods well known in the art. Methods of identifying suitable antigenic fragments of a protein are known in the art. See, *e.g.*, Hopp & Woods, 1981, *Proc. Natl. Acad. Sci. USA* 78:3824-3828; and

Jameson & Wolf, 1988, CABIOS (Computer Applications in the Biosciences) 4:181-186.

For the production of polyclonal antibodies, human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins or antigenic fragments, coupled to a suitable carrier, are injected on a periodic basis into an appropriate non-human host animal such as, *e.g.*, rabbits, sheep, goats, rats, mice or chickens. The animals are bled periodically (or eggs collected) and sera obtained are tested for the presence of antibodies to the injected subunit or antigen. The injections can be intramuscular, intraperitoneal, subcutaneous, and the like, and can be accompanied with adjuvant.

For the production of monoclonal antibodies, human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins or antigenic fragments, coupled to a suitable carrier, are injected into an appropriate non-human host animal as above for the production of polyclonal antibodies. In the case of monoclonal antibodies, the animal is generally a mouse. The animal's spleen cells are then immortalized, often by fusion with a myeloma cell, as described in Kohler & Milstein, 1975, Nature 256:495-497. For a fuller description of the production of monoclonal antibodies, see Antibodies: A Laboratory Manual, Harlow & Lane, eds., Cold Spring Harbor Laboratory Press, 1988.

Gene therapy may be used to introduce human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins into the cells of target organs. Nucleotides encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can be ligated into viral vectors which mediate transfer of the nucleotides by infection of recipient cells. Suitable viral vectors include retrovirus, adenovirus, adeno-associated virus, herpes virus, vaccinia virus, lentivirus, and polio virus based vectors. Alternatively, nucleotides encoding human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins can be transferred into cells for gene therapy by non-viral techniques including receptor-mediated targeted transfer using ligand-nucleotide conjugates, lipofection, membrane fusion, or direct microinjection. These procedures and variations thereof are suitable for *ex vivo* as well as *in vivo* gene therapy. Gene therapy with human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins will be particularly useful for the treatment of diseases where it is beneficial to elevate calcium sensitive potassium channel activity. cDNAs encoding mutant calcium

sensitive potassium channel subunits, that display a dominant negative phenotype, may be particularly useful for gene therapy treatment of diseases where it is beneficial to decrease calcium sensitive potassium channel activity.

The present invention includes processes for cloning orthologues of human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits from non-human species. In general, such processes include preparing a PCR primer or a hybridization probe based upon SEQ.ID.NO.:1, 3, 5, 7, 9, or 20 that can be used to amplify a fragment containing the non-human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit (in the case of PCR) from a suitable DNA preparation or to select a cDNA or genomic clone containing the non-human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit from a suitable library. A preferred embodiment of this process is a process for cloning the calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit from mouse.

By providing DNA encoding mouse calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits, the present invention allows for the generation of an animal model of human diseases in which calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit activity is abnormal. Such animal models can be generated by making transgenic "knockout" or "knockin" mice containing altered calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit genes. Knockout mice can be generated in which portions of the mouse calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit gene have been deleted. Knockin mice can be generated in which mutations that have been shown to lead to human disease are introduced into the mouse gene. Such knockout and knockin mice will be valuable tools in the study of the relationship between calcium sensitive potassium channels and disease and will provide important model systems in which to test potential pharmaceuticals or treatments for human diseases involving calcium sensitive potassium channels.

Accordingly, the present invention includes a method of producing a transgenic mouse comprising:

- (a) designing PCR primers or an oligonucleotide probe based upon SEQ.ID.NO.:1, 3, 5, 7, 9 or 20 for use in cloning the mouse calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit gene or cDNA;
- (b) using the PCR primers or the oligonucleotide probe to clone at least a portion of the mouse calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$,

or β 3d subunit gene or cDNA, the portion being large enough to use in making a transgenic mouse;

- (c) producing a transgenic mouse having at least one copy of the mouse calcium sensitive potassium channel β 2, β 3a, β 3b, β 3c, or β 3d subunit gene altered from its native state.

Methods of producing knockout and knockin mice are well known in the art. One method involves the use of gene-targeted ES cells in the generation of gene-targeted transgenic knockout mice and is described in, *e.g.*, Thomas et al., 1987, Cell 51:503-512, and is reviewed elsewhere (Frohman et al., 1989, Cell 56:145-147; Capecchi, 1989, Trends in Genet. 5:70-76; Baribault et al., 1989, Mol. Biol. Med. 6:481-492).

Techniques are available to inactivate or alter any genetic region to virtually any mutation desired by using targeted homologous recombination to insert specific changes into chromosomal genes. Generally, use is made of a "targeting vector," *i.e.*, a plasmid containing part of the genetic region it is desired to mutate. By virtue of the homology between this part of the genetic region on the plasmid and the corresponding genetic region on the chromosome, homologous recombination can be used to insert the plasmid into the genetic region, thus disrupting the genetic region. Usually, the targeting vector contains a selectable marker gene as well.

In comparison with homologous extrachromosomal recombination, which occurs at frequencies approaching 100%, homologous plasmid-chromosome recombination was originally reported to only be detected at frequencies between 10^{-6} and 10^{-3} (Lin et al., 1985, Proc. Natl. Acad. Sci. USA 82:1391-1395; Smithies et al., 1985, Nature 317: 230-234; Thomas et al., 1986, Cell 44:419-428). Nonhomologous plasmid-chromosome interactions are more frequent, occurring at levels 10^5 -fold (Lin et al., 1985, Proc. Natl. Acad. Sci. USA 82:1391-1395) to 10^2 -fold (Thomas et al., 1986, Cell 44:419-428) greater than comparable homologous insertion.

To overcome this low proportion of targeted recombination in murine ES cells, various strategies have been developed to detect or select rare homologous recombinants. One approach for detecting homologous alteration events uses the polymerase chain reaction (PCR) to screen pools of transformant cells for homologous insertion, followed by screening individual clones (Kim et al., 1988, Nucleic Acids Res. 16:8887-8903; Kim et al., 1991, Gene 103:227-233). Alternatively, a positive genetic selection approach has been developed in which a

marker gene is constructed which will only be active if homologous insertion occurs, allowing these recombinants to be selected directly (Sedivy et al., 1989, Proc. Natl. Acad. Sci. USA 86:227-231). One of the most powerful approaches developed for selecting homologous recombinants is the positive-negative selection (PNS) method
5 developed for genes for which no direct selection of the alteration exists (Mansour et al., 1988, Nature 336:348-352; Capecchi, 1989, Science 244:1288-1292; Capecchi, 1989, Trends in Genet. 5:70-76). The PNS method is more efficient for targeting genes which are not expressed at high levels because the marker gene has its own promoter. Nonhomologous recombinants are selected against by using the Herpes
10 Simplex virus thymidine kinase (HSV-TK) gene and selecting against its nonhomologous insertion with herpes drugs such as gancyclovir (GANC) or FIAU (1-(2-deoxy 2-fluoro-B-D-arabinofluranosyl)-5-iodouracil). By this counter-selection, the percentage of homologous recombinants in the surviving transformants can be increased.

15 Other methods of producing transgenic mice involve microinjecting the male pronuclei of fertilized eggs. Such methods are well known in the art.

The present invention includes a transgenic, non-human animal in which the animal's genome contains DNA encoding at least a portion of a human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit.

20

The following non-limiting examples are presented to better illustrate the invention.

EXAMPLE 1

25 Identification of the human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits and cDNA cloning

DNA sequence encoding the $\beta 1$ subunit was used to search the GenBank database for homologous sequences encoding novel subunits. This search yielded an EST with similarity to $\beta 1$ (AA904191). A cDNA encoding the EST was
30 purchased (Genome Systems) and sequenced in both directions. Synthetic oligonucleotide primers (SEQ.ID.NO.s:12 and 13) were used to amplify the coding region and a small portion of the 3' untranslated region (UTR) of this gene ($\beta 2$). The

coding region was then subcloned into a modified vector (pSP64T) containing an expanded polylinker between the 5' and 3' translation enhancer sequences (MVpl(+)).

The sequence of $\beta 2$ was then used to search the GenBank database for additional novel beta subunits. The sequences from identified EST's were then used
5 to search the database again. Several EST's were obtained in this iterative approach: AA195381, AA236930, AA236968, AA279911, AA761761, AA934876, AA195511, AA917510. The alignment of these sequences suggested they encoded the C-terminal portion of a novel β subunit, here designated $\beta 3$. Available cDNAs encoding these
10 ESTs were purchased (Genome Systems) and sequenced in both directions. None of these clones encoded full length protein based on the lack of 5' in-frame stop codons and amino acid alignments only to the middle of the first transmembrane segments of $\beta 1$ and $\beta 2$.

Unique and conserved portions of the individual subunits were used separately to search the databases for genomic sequences encoding these transcripts.
15 A single 180 kilobase fragment of unidentified genomic sequence was identified using $\beta 3a$, $\beta 3b$ and $\beta 3c$ specific fragments (GenBank accession number AC007823, version 2). Later versions of this entry contained a 40.4 kilobase contiguous fragment that contained all three specific fragments in the following order $\beta 3a$, $\beta 3b$ and $\beta 3c$. $\beta 3c$ is contiguous with the 5' end of the core sequence. See Figure 8.

20 A synthetic oligo, 5'-TTT ACA TTG TTA GTT TGC AGA CAG G-3' (SEQ.ID.NO.:19), annealing 3' of the $\beta 3$ stop codon was used in a 5' RACE reaction as described in Clontech's Marathon Ready Spleen cDNA kit (catalog # 7412-1). This reaction yielded multiple products of varying sizes. Several fragments separated by electrophoresis were extracted from gel slices and cloned. Three distinct subunits
25 were identified ($\beta 3a$, $\beta 3b$ and $\beta 3c$) in this manner.

To ensure novel subunits were not overlooked, the unfractionated product of the PCR amplification reaction was cloned directly into a TA cloning vector (pCR2.1, Invitrogen), without any attempt to isolate specific fragments. Colonies were then screened using a probe derived from EST AA761761 by the
30 'colony filter hybridization protocol' as described in *Current Protocols in Molecular Biology*, sections 6.1.1 and 6.3.1. DNA was prepared from hybridizing colonies. cDNAs with restriction digest patterns distinct from the original clones were sequenced in both directions. The open reading frames were determined and

amplified using synthetic oligonucleotide primers (SEQ.ID.NOs.:14 through 18), and subcloned into MVpl(+). One additional unique subunit was identified: β 3d.

EXAMPLE 2

5 Analysis of expression of human calcium sensitive potassium channel β 2, β 3a, β 3b, β 3c, or β 3d subunits

Northern blot analysis: Northern blots containing poly(A+)-RNA from human heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testes, ovary, small intestine, colon, and peripheral blood
 10 leukocytes were purchased from Clontech, Palo Alto, CA. The blots were probed with ^{32}P -labeled, randomly primed cDNA probes from β 2 (nucleotides 268 to 1080 of SEQ.ID.NO.:1), β 3a (nucleotides 70 to 384 of SEQ.ID.NO.:3), β 3b (nucleotides 463 to 797 of SEQ.ID.NO.:5), and β 3c/d (nucleotides 311 to 912 of SEQ.ID.NO.:7). The hybridization was carried out in 5X SSPE, 10X Denhardts solution, 50%
 15 Formamide, 2% SDS, 100ug/ml salmon sperm DNA at 42°C overnight. The washes were carried out stepwise in 2X SSC, 0.05% SDS at 42°C for 40 minutes, followed by 1X SSC, 0.05% SDS at 50°C for 40 minutes. High stringency washes were carried out at 0.1SSC, 0.05% SDS at 65°C for 40 minutes. Hybridization was detected either by exposure of the washed blots to X-ray film or by electronic detection using a
 20 phosphorimager.

Electrophysiological analysis: cRNAs were synthesized in vitro from plasmids encoding human *Slowpoke* α or the β 2, β 3a, β 3b, β 3c, or β 3d subunits and injected into *Xenopus* oocytes (1.5 ng/oocyte of α subunit RNA +/- β subunit RNA at 1, 5, or 10X molar excess). Calcium sensitive potassium currents were recorded in
 25 inside-out patches. Recordings were performed under ionic conditions of symmetrical potassium. The standard pipette and bath solutions contained 116 mM potassium gluconate, 4 mM potassium chloride, 10 mM HEPES, pH 7.2. CaCl_2 was added to the bath solution to give final concentrations of free ionized calcium of 3-30 μM , taking into account the stability constant for calcium gluconate (15.9 M^{-1}).
 30 Currents were recorded using an EPC-7 amplifier (HEKA). The pClamp6.0 program (Axon Instruments) was used to generate voltage-clamp commands for data acquisition, and for analysis. NP_o - voltage relations were determined at 3, 10 and 30

μM bath calcium using two methods: (1) calculation of macroscopic conductance from peak or steady-state currents at test potentials (-80 to 80 mV), or (2) measurement or calculation of tail currents peaks (-80 mV) at test potentials. Boltzmann functions were fit to the data and used to derive the half-maximal activation parameter ($V_{1/2}$). Maximal inactivation parameters (30 μM Ca^{2+} and 80 mV) were calculated from current traces or averaged current traces. Inactivation rates were determined from single exponential fits. Fractional non-inactivating current was calculated as steady-state/peak current; fractional inactivating current was estimated as peak current minus steady-state current divided by peak current.

10

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims.

15

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

WHAT IS CLAIMED IS:

1. An isolated DNA comprising nucleotides encoding a human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein.
5
2. The DNA of claim 1 comprising nucleotides encoding a polypeptide having an amino acid sequence selected from the group consisting of: SEQ.ID.NO.:2; SEQ.ID.NO.:4; SEQ.ID.NO.:4 with an asparagine at position 163 instead of a serine; SEQ.ID.NO.:6; SEQ.ID.NO.:6 with a serine at position 143
10 instead of an asparagine; SEQ.ID.NO.:8; SEQ.ID.NO.:8 with an asparagine at position 161 instead of a serine; SEQ.ID.NO.:10; SEQ.ID.NO.:10 with a serine at position 165 instead of an asparagine; and positions 2-246 of SEQ.ID.NO.:6.
3. The DNA of claim 1 comprising a nucleotide sequence selected
15 from the group consisting of: SEQ.ID.NO.:1, 3, 5, 7, 9, and 20.
4. The DNA of claim 1 comprising a nucleotide sequence selected from the group consisting of: positions 271-975 of SEQ.ID.NO.:1, positions 341 to 1171 of SEQ.ID.NO.:3, positions 796 to 1566 of SEQ.ID.NO.:5, positions 869 to
20 1693 of SEQ.ID.NO.:7, and positions 457 to 1293 of SEQ.ID.NO.:9.
5. An isolated DNA that hybridizes under stringent conditions to the DNA of claim 2.
- 25 6. An expression vector comprising the DNA of claim 1.
7. A recombinant host cell comprising the DNA of claim 1.
8. An isolated human calcium sensitive potassium channel $\beta 2$,
30 $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein.
9. The protein of claim 8 having an amino acid sequence selected from the group consisting of: SEQ.ID.NO.:2; SEQ.ID.NO.:4; SEQ.ID.NO.:4 with an asparagine at position 163 instead of a serine; SEQ.ID.NO.:6; SEQ.ID.NO.:6 with a

serine at position 143 instead of an asparagine; SEQ.ID.NO.:8; SEQ.ID.NO.:8 with an asparagine at position 161 instead of a serine; SEQ.ID.NO.:10; SEQ.ID.NO.:10 with a serine at position 165 instead of an asparagine; and positions 2-246 of SEQ.ID.NO.:6.

- 5 10. The protein of claim 8 containing a single amino acid substitution.
11. The protein of claim 8 containing two or more amino acid substitutions where the amino acid substitutions do not occur in conserved positions.
- 10 12. A polypeptide having at least 80% sequence identity to the protein of claim 9 when measured by BLAST or FASTA.
13. An antibody that binds specifically to a human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit protein; or that binds specifically to the $\beta 3$ subunit family of proteins by binding to the conserved core.
- 15 14. A DNA or RNA oligonucleotide probe comprising at least 15 contiguous nucleotides of at least one of a sequence selected from the group consisting of: SEQ.ID.NO.:1, 3, 5, 7, 9, and 20.
- 20 15. A method for identifying substances that bind to calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins comprising:
- 25 (a) providing cells expressing a calcium sensitive potassium channel containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;
- (b) exposing the cells to a substance that is not known to bind calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;
- 30 (c) determining the amount of binding of the substance to the cells;
- (d) comparing the amount of binding in step (c) to the amount of binding of the substance to control cells where the control cells are substantially

identical to the cells of step (a) except that the control cells do not express human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

where if the amount of binding in step (c) is greater than the amount of binding of the substance to control cells, then the substance binds to calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

16. A method of identifying substances that bind calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins and thus are likely to be inhibitors or activators of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins comprising:

(a) providing cells expressing calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

(b) exposing the cells to a compound that is known to bind to the calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

(c) determining the amount of binding of the compound to the cells in the presence and in the absence of a substance not known to bind to calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

where if the amount of binding of the compound in the presence of the substance differs from that in the absence of the substance, then the substance binds calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins and is likely to be an inhibitor or activator of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

17. A method of identifying activators or inhibitors of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins comprising:

(a) recombinantly expressing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins or mutant human calcium

sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins in a host cell so that the recombinantly expressed human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins form calcium sensitive potassium channels by forming heteromers with other calcium sensitive potassium channel subunit proteins;

5 (b) measuring the biological activity of the calcium sensitive potassium channels formed in step (a) in the presence and in the absence of a substance suspected of being an activator or an inhibitor of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins;

10 where a change in the biological activity of the calcium sensitive potassium channels formed in step (a) in the presence as compared to the absence of the substance indicates that the substance is an activator or an inhibitor of calcium sensitive potassium channels containing human calcium sensitive potassium channel $\beta 2$, $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunit proteins.

15

18. A method of identifying DNA sequences in the $\beta 3$ gene that promote, enhance, or repress gene transcription comprising:

(a) constructing a promoter-reporter vector such that fragments of the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) precede the coding cDNA sequence of a reporter gene which encodes a reporter protein;

20

(b) transfecting the vector into cells and measuring the abundance of the reporter protein encoded by the vector;

(c) comparing the abundance of the reporter protein in the cells of step (b) to the abundance of the reporter protein in cells transfected with the vector without fragments of the promoter region of the $\beta 3$ gene;

25

where fragments of the promoter region of the $\beta 3$ gene which increase the abundance of the reporter protein in the absence of other promoter elements only in cells which endogenously express $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits are promoter elements; sequences which decrease the abundance of the reporter protein in the presence of an unrelated constitutive promoter element in cells which do not endogenously express $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits are repressor elements; and sequences which increase the abundance of the reporter protein in the presence of an unrelated constitutive promoter element in cells which endogenously express $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits are enhancer elements.

30

19. The method of claim 18 where the vector contains promoter or enhancer sequence elements which function independently of the fragments of the promoter region of the $\beta 3$ gene.

5

20. The method of claim 18 where the abundance of the reporter protein is normalized with respect to the fraction of transfected cells.

21. A method of identifying DNA sequences in the $\beta 3$ gene that promote, enhance, or repress gene transcription comprising:

10

(a) incubating radiolabeled fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) with nuclear extracts from cells; and

15

(b) separating the incubation on a gel; where fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene that migrate differently in a gel ('undergo a shift') after incubation with nuclear extracts from cells are DNA sequences which bind nuclear factors which promote, enhance or repress $\beta 3$ gene expression.

20

22. The method of claim 21 where the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the method of claim 18.

25

23. The method of claim 21 where the cells express $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits.

24. The method of claim 21 where the cells do not express $\beta 3a$, $\beta 3b$, $\beta 3c$, or $\beta 3d$ subunits.

30

25. A method of identifying nuclear factors involved in $\beta 3$ gene transcription regulation comprising:

(a) incubating radiolabeled fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene

(SEQ.ID.NO.:20, nucleotides 1 to 17,436) with cloned or purified transcription factors and separating the incubation on a gel;

5 where factors which bind $\beta 3$ gene promoter sequence elements will induce a shift in the migration of the radiolabeled DNA fragments, and are involved in $\beta 3$ gene transcription regulation.

10 26. The method of claim 25 where the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

27. A method of identifying transcription factors involved in $\beta 3$ gene transcription regulation comprising:

15 (a) incubating radiolabeled fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) with nuclear extracts from cells and separating the incubation on a gel;

(b) adding an antibody that specifically recognizes a single transcription factor or a family of transcription factors to the incubation of step (a), followed by separating the incubation on a gel;

20 where a super-shift in mobility of the double stranded DNA in step (b) as compared to step (a) indicates that a transcription factor recognized by the antibody binds the double stranded DNA.

25 28. A method of identifying clones encoding nuclear factors involved in $\beta 3$ gene transcription regulation by cloning comprising:

(a) screening an expression library with radiolabeled fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436)

30 (b) determining which clones of the library bind the radiolabeled fragments of double stranded DNA;

(c) amplifying and sequencing the clones of step (b).

29. The method of claim 28 where the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

5 30. A method of identifying nuclear factors involved in $\beta 3$ gene transcription regulation by cloning comprising:

(a) attaching fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) to a stable matrix;

10 (b) incubating phage expressing cDNA encoded fusion proteins at their surface with the matrix;

(c) removing phage that do not bind to the matrix by washing;

(d) eluting phage bound to the matrix with excess fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene;

15 where the phage eluted in step (d) encode nuclear factors involved in $\beta 3$ gene transcription regulation.

20 31. The method of claim 30 where the DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

32. The method of claim 30 where the phage eluted at step (d) are amplified and sequenced.

25

33. A method of identifying nuclear factors involved in $\beta 3$ gene transcription regulation comprising:

(a) attaching fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) to a stable matrix;

30

(b) incubating nuclear extracts from cells with the matrix;

(c) washing non-binding proteins from the nuclear extract from the matrix;

(d) eluting bound proteins from the matrix with excess double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene;

5 where the eluted proteins from step (d) are nuclear factors involved in $\beta 3$ gene transcription regulation.

34. The method of claim 33 further comprising separating the eluted proteins from step (d) on a gel and staining the gel to test for purity of the eluted proteins.

10

35. The method of claim 34 further comprising sequencing the proteins that have been separated on the gel.

15 36. The method of claim 34 further comprising immunological analysis of the proteins that have been separated on the gel with antibodies directed towards known transcription factors to identify the eluted proteins by western blot or immunoprecipitation.

20 37. The method of claim 33 where the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

38. A method of identifying nuclear factors involved in $\beta 3$ gene transcription regulation by cloning comprising:

25 (a) constructing a yeast strain that contains a few to several copies of a fragment of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) preceding a cDNA encoding a reporter protein;

30 (b) constructing a cDNA library from cells in a vector that allows formation of fusion proteins encoded by the inserted cDNA and a transcription activation domain;

(c) transforming the library of (b) into the yeast strain of (a) and isolating colonies of yeast displaying expression of the reporter protein.

39. The method of claim 38 where the fragments of double stranded DNA corresponding to sequences found in the promoter region of the $\beta 3$ gene are identified by the methods of claim 18 or 21.

5 40. The method of claim 38 further comprising purifying the vectors from the isolated colonies and sequencing the cDNA in the vectors.

41. A method of identifying substances that enhance or inhibit the rate of transcription of the $\beta 3$ gene comprising:

- 10 (a) constructing a promoter-reporter vector such that fragments of the promoter region of the $\beta 3$ gene (SEQ.ID.NO.:20, nucleotides 1 to 17,436) precede the coding cDNA sequence of a reporter gene which encodes a reporter protein;
- (b) transfecting the vector into cells and measuring the abundance of the reporter protein encoded by the vector in the presence and absence of a
- 15 compound;

 where (1) if the presence of the compound decreases the abundance of the reporter protein, then the compound is a substance that inhibits the rate of transcription of the $\beta 3$ gene; (2) if the presence of the compound increases the abundance of the reporter protein, then the compound is a substance that enhances the

20 rate of transcription of the $\beta 3$ gene.

42. The method of claim 41 further comprising a control in which the effect of the compound on the abundance of the reporter protein in control cells is measured, where the control cells are cells that are essentially the same as the cells of

25 step (b) except that the control cells have been transfected with a vector that lacks fragments of the promoter region of the $\beta 3$ gene.

FIGURE 1A

```
1  CTTAATCCTA  TCCAAGTATG  CAGTACGCTC  TTGGGTCGTC  TCATGAGACC  CAGGGGCATG
61  TTGGAAGAA  CTGAGAGAAA  GAGCAACAAA  GCGGCGAGTG  GTGTGAGAGG  GCAGCACGCG
121 CTGTGGGGCC  CTTCCAGAGA  AATGTACTGA  AAAAGTCTAC  GCAATGTCTG  GGATTTGCTA
181 AACAAATACCT  GGAAAGCAGA  CAGGTTTTTT  TGCCATTCCCT  CCAGGACATC  CACCATAAGG
241 AAAGGAGACC  CTGGACCAAC  ATTCTCTAAG  ATGTTTATAT  GGACCAGTGG  CCGGACCTCT
301 TCATCTTATA  GACATGATGA  AAAAAGAAAT  ATTTACCAGA  AAATCAGGGA  CCATGACCTC
361 CTGGACAAAA  GGAAAACAGT  CACAGCACTG  AAGGCAGGAG  AGGACCGAGC  TATTCTCCTG
421 GGACTGGCTA  TGATGGTGTG  CTCCATCATG  ATGTATTTTC  TGCTGGGAAT  CACACTCCTG
481 CGCTCATACA  TGCAGAGCGT  GTGGACCGAA  GAGTCTCAAT  GCACCTTGCT  GAATGCGTCC
541 ATCACGGAAG  CATTTAACTG  CTCCTTCAGC  TGTGGTCCAG  ACTGCTGGAA  ACTTTCTCAG
601 TACCCCTGCC  TCCAGGTGTA  CGTTAACCTG  ACTTCTTCCG  GGGAAAAGCT  CCTCCTCTAC
661 CACACAGAAG  AGACAATAAA  AATCAATCAG  AAGTGCTCCT  ATATACCTAA  ATGTGGAAAA
721 AATTTTGAAG  AATCCATGTC  CCTGGTGAAT  GTTGTCTATG  AAAACTTCAG  GAAGTATCAA
781 CACTTCTCCT  GCTATTCTGA  CCCAGAAGGA  AACCAGAAGA  GTGTTATCCT  AACCAAACTC
841 TACAGTTCCA  ACGTGCTGTT  CCATTCACCT  TTCTGGCCAA  CCTGTATGAT  GGCTGGGGGT
901 GTGGCAATTG  TTGCCATGGT  GAAACTTACA  CAGTACCTCT  CCCTACTATG  TGAGAGGATC
961 CAACGGATCA  ATAGATAAAT  GCAAAAATGG  ATAAAATAAT  TTTTGTTAAA  GCTCAAATAC
1021 TGTTTTCTTT  CATTCTTCAC  CAAAGAACCT  TAAGTTTGTA  ACGTGCAGTC  TGTATGAGT
1081 TCCCTAATAT  ATTCTTATAT  GTAGAGCAAT  AATGCAAAAG  CTGTTCTATA  TGCAAACATG
1141 ATGTCTTTAT  TATTCAGGAG  AATAAATAAC  TGTTTTGTGT  TGAA
```

FIGURE 1B

1 MFIWTSGRTS SSYRHDEKRN IYQKIRDHDL LDKRKTVTAL KAGEDRAILL GLAMMVCSIM
61 MYFLLGITLL RSYMQSVWTE ESQCTLLNAS ITETFNCFSFS CGPDCWKLSQ YPCLQVYVNL
121 TSSGEKLLLY HTEETIKINQ KCSYIPKCGK NFEESSMLVN VVMENFRKYQ HFSCYSDPEG
181 NQKSVILTKL YSSNVLFHSL FWPTCMMAGG VAIVAMVKLT QYLSLLCERI QRINR

FIGURE 2A

```

1  GCTCCCGGCT  GCCGAGGCGG  AAACACAGGT  GATGAGGTGG  CGGCAAGCAC  AGTGCAAAGA
61  GAGAGAAGCA  GCTTCGGCTG  CAGCAAACCA  CGCAGGTCCT  TCTTGATCAT  CTAGAACTGA
121  CCGCTCCGCC  TTGCCAGGAG  TCTGCAGAAC  CACGTGGCTG  GCCTGCCTGA  AGTTCTCACC
181  TCTCTAGGAA  GCGGGGGGGC  TTCTAATGGC  TGCAGCTGCG  CTGGGGGGCTG  GGGGCTCCCG
241  CTGGGACTCC  ACTTCCGTGG  ATGTCTAAGC  TTCACCTTTC  TTGCGCCCGC  AGGGGCATGA
301  CTCAGGTGAA  AGGGAGCCAT  TTTCTCAGAC  CCCTGGCCTC  ATGCAGCCCT  TCAGCATCCC
361  CGTGCAAATC  AACTTCAGG  GCAGCCGGAG  GCGCCAGGGG  AGGACAGCCT  TTCCTGCCTC
421  AGGGAAGAAG  AGAGAGACAG  ACTACAGTGA  TGGAGACCCA  CTAGATGTGC  ACAAGAGGCT
481  GCCATCCAGT  ACTGGAGAGG  ACCGAGCCGT  GATGCTGGGG  TTTGCCATGA  TGGGCTTCTC
541  AGTCCTAATG  TTCTTCTTGC  TCGGAACAAC  CATTCTAAAG  CCTTTTATGC  TCAGCATTCA
601  GAGAGAAGAA  TCGACCTGCA  CTGCCATCCA  CACAGATATC  ATGGACGACT  GGCTGGACTG
661  TGCCTTCACC  TGTGGTGTGC  ACTGCCACGG  TCAGGGGAAG  TACCCGTGTC  TTCAGGTGTT
721  TGTGAACCTC  AGCCATCCAG  GTCAGAAAGC  TCTCCTACAT  TATAATGAAG  AGGCTGTCCA
781  GATAAATCCC  AAGTGCTTTT  ACACACCTAA  GTGCCACCAA  GATAGAAGTG  ATTTGCTCAA
841  CAGTGCTCTG  GACATAAAAG  AATTCTTCGA  TCACAAAAAT  GGAACCCCTT  TTTCATGCTT
901  CTACAGTCCA  GCCAGCCAAT  CTGAAGATGT  CATTCTTATA  AAAAAGTATG  ACCAAATGGC
961  TATCTTCCAC  TGTTTATTTT  GGCCCTCACT  GACTCTGCTA  GGTGGTGCCC  TGATTGTTGG
1021  CATGGTGAGA  TTAACACAAC  ACCTGTCCTT  ACTGTGTGAA  AAATATAGCA  CTGTAGTCAG
1081  AGATGAGGTA  GGTGGAAAAG  TACCTTATAT  AGAACAGCAT  CAGTTCAAAC  TGTGCATTAT
1141  GAGGAGGAGC  AAAGGAAGAG  CAGAGAAATC  TTAAGACGGT  GGCCAAATTA  AAGTGCTGGC
1201  CTTCAGATGT  CTGTGATTTT  TGCAACTCGA  GTATGCG

```


FIGURE 2B

1 MQPFSIPVQI TLQGSRRRQG RTAFPASGKK RETDYSDDGP LDVHKRLPSS TGEDRAVMLG
61 FAMMGFSVLM FLLGTTILK PFMLSIQREE STCTAIHTDI MDDWLDCAFT CGVHCHGQGK
121 YPCLQVFNVL SHPGQKALLH YNEEAVQINP KCFYTPKCHQ DRSDLLNSAL DIKEFFDHKN
181 GTPFSCFYSP ASQSEDVILI KKYDQMAIFH CLFWPSLTLL GGALIVGMVR LTQHLSLLCE
241 KYSTVVRDEV GGVVPYIEQH QFKLCIMRRS KGRAEKS

FIGURE 3A

```

1  AAGAGAAAGA ACAAGAAAAA GAAAAAGAAG AGGAAAAAAT CCCCAGTACC CATAGAAACC
61  CTTAAAGATG TTTAAAAAGA GTTAACTTAT CAGAACACAG ATTTAAGTGA AATTAAGGAA
121 GAAGAGCAGG TAAAGTCTAC TGACAGAAAG TCAGCAGTGG AAGCCCAAAA CGAGGTGACT
181 GAAAATCCAA AACAGAAAAAT TGCAGCAGAA AGCAGTGAAA ATGCTGATTG TCCAGAGAAT
241 CCTAAATGA AGTTGGATGG AAAACTTGAC CAAGAAGGCA ATGATGTAAA AACAGCAGCT
301 GAGGAGGTAC TAGCTGGTAG AGACACATTA GATTTTGAGG ATGTCACAGT TCAATCATCA
361 GGCCCGAGGG CTGGTGGTGA AGAATTAGAT GAAGGTGTTG CAAAAGATAA TGCTAAAATA
421 GCTGGTGCCA CTTAAAGCAA TCCTGAAGAA CCAGAGAGTG AAGATGCAGA TCACTGCACC
481 GTACCCAAAA ATGAAAGTCC CTCACAGGAC ATTAGTGATG CCTGTGAAGC AGAAAGTACA
541 GAGAGGTGCG GGATGTCAGA ACATCCAAGT CAGACCATCA GGAAAGCTTT AGACAGCAAT
601 AGCCTAAAAA ACCATGACTT GTTGGCACCA GGAGGAGAGC CGGGGGACTT CAATCCAGAA
661 AGCAGAGAAG ATACCAGAGG AGGGAACGAG AAGGGCAAAA GCAAAGAAGA CCGTACCATG
721 TCCTAAGCTG AGGCAGGCGG CAGGCGTGGT GCACAGGAAG TCTGAGTGTG AGGGGCTCTT
781 TTCTCTCCAC TGCCAATGAC AGCCTTTTCCT GCCTCAGGGA AGAAGAGAGA GACAGACTAC
841 AGTGATGGAG ACCCACTAGA TGTGCACAAG AGGCTGCCAT CCAGTACTGG AGAGGACCGA
901 GCCGTGATGC TGGGGTTTGC CATGATGGGC TTCTCAGTCC TAATGTTCTT CTTGCTCGGA
961 ACAACCATTC TAAAGCCTTT TATGCTCAGC ATTCAGAGAG AAGAATCGAC CTGCACTGCC
1021 ATCCACACAG ATATCATGGA CGACTGGCTG GACTGTGCCT TCACCTGTGG TGTGCAATGC
1081 CACGGTCAGG GGAAGTACCC GTGTCTTCAG GTGTTTGTGA ACCTCAGCCA TCCAGGTCAG
1141 AAAGTCTCC TACATTATAA TGAAGAGGCT GTCCAGATAA ATCCCAAGTG CTTTACACA
1201 CCTAAGTGCC ACCAAGATAG AAATGATTTG CTCAACAGTG CTCTGGACAT AAAAGAATTC
1261 TTCGATCACA AAAATGGAAC CCCCTTTTCA TGCTTCTACA GTCCAGCCAG CCAATCTGAA
1321 GATGTCATTC TTATAAAAAA GTATGACCAA ATGGCTATCT TCCACTGTTT ATTTTGGCCT
1381 TCACTGACTC TGCTAGGTGG TGCCCTGATT GTTGGCATGG TGAGATTAA ACACACCTG
1441 TCCTTACTGT GTGAAAAATA TAGCACTGTA GTCAGAGATG AGGTAGGTGG AAAAGTACCT
1501 TATATAGAAC AGCATCAGTT CAAACTGTGC ATTATGAGGA GGAGCAAAGG AAGAGCAGAG
1561 AAATCTTAAG ACGGTGGCCA AATTAAAGTG CTGGCCTTCA GATGTCTGTG ATTTCTGCAA
1621 CTCGAGTATG CG

```

FIGURE 3B

1 MTAFPASGKK RETDYSDBGD LDVHKRLPSS TGEDRAVMLG FAMMGFSVLM FLLGTTILK
61 PFMLSIQREE STCTAIHTDI MDDWLDCAFT CGVHCHGQ GK YPCLQVFVNL SHPGQKALLH
121 YNEEAVQINP KCFYTPKCHQ DRNDLLNSAL DIKEFFDHKN GTPFSCFYSP ASQSEDVILI
181 KKYDQMAIFH CLFWPSLTLL GGALIVGMVR LTQHLSLLCE KYSTVVRDEV GGKVPYIEQH
241 QFKLCIMRRS KGRAEKS

FIGURE 4A

```

1 CCCAGCTACT CGGGAGGCTG AGGCAGGAGA ATCGCTTGAA CCTGGGAGGC GGAGGAGGTT
61 GCAGTGAAC T GAGATCGTAC CCAGCCTGGG CAACAGTGCG AGGCTCCGTC TCAAAAAAAA
121 ACCAAAAAAC ACAAAAACAA AAAACGACAG AGAAGGCCAA ACAAAACACA TCTGTGGGCT
181 GGATGCCGCC ATGCCCACCG GTTTGCGACC TTTGTGTTGG ACTCTTCTGT TCACCAGACA
241 CCCTGCCCTG CGAGAATGTA TCTCATCCTT TGCTGGAGCA GGTTCGAGG CACAGTGGAG
301 AGAGGAGAGA AGAAATGAAG GGACACTTAT GCAGAACCAT GAGTGGCCAG AGAGGAGGAG
361 AAGGAGGGTG AGAGGAGCAA AGAAGCCATG ACAACTTCAT AATTCTGAGT GGACTGGGCA
421 GTGGCCAGAA ATTCTGGTGG TGGATATGCT GCCTTTCCAA CAGGTGAATA TGAAAGAATA
481 AGTCAAACCC TGTTCAGGAC GCTGTTAATT CCAAATGTGA ACTTTTGTAG TCATTCTTTT
541 CATGTGGAAT TCAAAGGAGA ATGTAAACAA ATTTTCAGGA GGGACGTGCA ATATCCCTGA
601 AAGATAACAG AGTTCGTAA CACTTATTTAC ATACAACATT CTCTAGTTAT TGATTAAACA
661 GATCTCTACA GACTTGCATG AGGCAACATT TCTTAGGCTT GTTTGCTACA ATATCTTTAA
721 AAATACTTGA TTACACATCA CTTTAGCTTA TTTAGATGGA CTTTTCACCA AGCTCTGAAC
781 TGGGATTTCA TTTTGTGTC TTTTATCCTGC TCACGAGACA CAGGTAGGCA GCAAATGAGA
841 TTATCCCTCC AGTCCCCATG GATTGGAAAT GTTCCCCCTT CTTTATGAGC TCACTGCAGT
901 ATCTCCTTCT CCCTTTCCCC AAAGGACAGC CTTTCCTGCC TCAGGGAAGA AGAGAGAGAC
961 AGACTACAGT GATGGAGACC CACTAGATGT GCACAAGAGG CTGCCATCCA GTACTGGAGA
1021 GGACCGAGCC GTGATGCTGG GGTTCGCCAT GATGGGCTTC TCAGTCCTAA TGTTCCTCTT
1081 GCTCGGAACA ACCATTCTAA AGCCTTTTAT GCTCAGCATT CAGAGAGAAG AATCGACCTG
1141 CACTGCCATC CACACAGATA TCATGGACGA CTGGCTGGAC TGTGCCTTCA CCTGTGGTGT
1201 GCACTACCAC GGTCAGGGGA AGTACCCGTG TCTTCAGGTG TTTGTGAACC TCAGCCATCC
1261 AGGTCAGAAA GCTCTCCTAC ATTATAATGA AGAGGCTGTC CAGATAAATC CCAAGTGCTT
1321 TTACACACCT AAGTGCCACC AAGATAGAAG TGATTTGCTC AACAGTGCTC TGGACATAAA
1381 AGAATTCTTC GATCACAAAA ATGGAACCCC CTTTTCATGC TTCTACAGTC CAGCCAGCCA
1441 ATCTGAAGAT GTCATTCTTA TAAAAAAGTA TGACCAAATG GCTATCTTCC ACTGTTTATT
1501 TTGGCCTTCA CTGACTCTGC TAGGTGGTGC CCTGATTGTT GGCATGGTGA GATTAAACACA
1561 ACACCTGTCC TTACTGTGTG AAAAATATAG CACTGTAGTC AGAGATGAGG TAGGTGGAAA
1621 AGTACCTTAT ATAGAACAGC ATCAGTTCAA ACTGTGCATT ATGAGGAGGA GCAAAGGAAG
1681 AGCAGAGAAA TCTTAAGACG GTGGCCAAAT TAAAGTGCTG GCCTTCAGAT GTCTGTGATT
1741 TCTGCAACTC GAGTATGCG

```

FIGURE 4B

1 MFPLLYELTA VSPSPFPQRT AFPASGKKRE TDYSDGDPLD VHKRLPSSTG EDRAVMLGFA
61 MMGFSVLMFF LLGTTILKPF MLSIQREEST CTAIHTDIMD DWLDCAFTCG VHCHGQGKYP
121 CLQVFVNLSH PGQKALLHYN EEAVQINPKC FYTPKCHQDR SDLLNSALDI KEFFDHKNGT
181 PFSCFYSPAS QSEDVILIKK YDQMAIFHCL FWPSLTLLGG ALIVGMVRLT QHLSLLCEKY
241 STVVRDEVGG KVPYIEQHOF KLCIMRRSKG RAEKS

FIGURE 5A

```

1  CGCCGCGGAT CCGAAATGAA GGGACACTTA TGCAGAACCA TGAGTGGCCA GAGAGGAGGA
61 GAAGGAGGGT GAGAGGAGCA AAGAAGCCAT GACAACTTCA TAATTCTGAG TGGACTGGGC
121 AGTGGCCAGA AATTCTGGTG GTGGATATGC TGCCTTTCCA ACAGGTGAAT ATGAAAGAAT
181 AAGTCAAACC CTGTTTCAGGA CGCTGTAAAT TCCAAATGTG AACTTTTTGA GTCATTCTTT
241 TCATGTGGAA TTCAAAGGAG AATGTAAACA AATTTTCAGG AGGGACGTGC AATATCCCTG
301 AAAGATAACA AAGTTCGTAA CACTTATTTA CATAAACAT TCTCTAGTTA TTGATTAAAC
361 AGATCTCTAC AGACTTGCAT GAGGCAACAT TTCTTAGGCT TGTTTGCTAC AATATCTTTA
421 AAAATACTTG ATTACACATC ACTTTAGCTT ATTTAGATGG ACTTTTCACC AAGCTCTGAA
481 CTGGGATTTT ATTTTGTTGC ATTCATCCTG CTCACGAGAC ACAGGACAGC CTTTCCTGCC
541 TCAGGGAAGA AGAGAGAGAC AGACTACAGT GATGGAGACC CACTAGATGT GCACAAGAGG
601 CTGCCATCCA GTACTGGAGA GGACCGAGCC GTGATGCTGG GGTTCGCCAT GATGGGCTTC
661 TCAGTCCTAA TGTTCTTCTT GCTCGGAACA ACCATTCTAA AGCCTTTTAT GCTCAGCATT
721 CAGAGAGAAG AATCGACCTG CACTGCCATC CACACAGATA TCATGGACGA CTGGCTGGAC
781 TGTGCCTTCA CCTGTGGTGT GCACTGCCAC GGTGAGGGA AGTACCCGTG TCTTCAGGTG
841 TTTGTGAACC TCAGCCATCC AGGTCAGAAA GCTCTCCTAC ATTATAATGA AGAGGCTGTC
901 CAGATAAATC CCAAGTGCTT TTACACACCT AAGTGCCACC AAGATAGAAA TGATTTGCTC
961 AACAGTGCTC TGGACATAAA AGAATTCTTC GATCACAAAA ATGGAACCCC CTTTTCATGC
1021 TTCTACAGTC CAGCCAGCCA ATCTGAAGAT GTCATTCTTA TAAAAAAGTA TGACCAAATG
1081 GCTATCTTCC ACTGTTTATT TTGGCCTTCA CTGACTCTGC TAGGTGGTGC CCTGATTGTT
1141 GGCATGGTGA GATTAACACA ACACCTGTCC TTACTGTGTG AAAAATATAG CACTGTAGTC
1201 AGAGATGAGG TAGGTGGAAG AGTACCTTAT ATAGAACAGC ATCAGTTCAA ACTGTGCATT
1261 ATGAGGAGGA GCAAAGGAAG AGCAGAGAAA TCTTAA

```

FIGURE 5B

1 MDFSPSSELG FHFVAFILLT RHRTAFPASG KKRETDYSDG DPLDVHKRLP SSTGEDRAVM
61 LGFAMMGFSV LMFFLLGTI LKPFMLSIQR EESTCTAIHT DIMDDWLDCA FTCGVHCHGQ
121 GKYPCLQVFV NLSHPGQKAL LHYNEEAVQI NPKCFYTPKC HQDRNDLLNS ALDIKEFFDH
181 KNGTPFSCFY SPASQSEDVI LIKKYDQMAI FHCLFWPSLT LLGGALIVGM VRLTQHLSLL
241 CEKYSTVVVD EVGGKVPYIE QHQFKLCIMR RSKGRAEKS

FIGURE 6

```

BKb1 .....MVK.KLVM
BKb2 .....MFIWTSGR TSSSYRHDEKRNIYQKIRDHDL LDKRKTVT
BKb3a .....MQPFSIPVQITLQGSRRRQGR TAFPASGKKRETDYS...DGDPLDVHKRLP
BKb3b .....MTAFPASGKKRETDYS...DGDPLDVHKRLP
BKb3c .....MFPLLYELTAVSPSPFPQRTAFPASGKKRETDYS...DGDPLDVHKRLP
BKb3d MDFSPSELGFHFVAFILLTRH.....RTAFPASGKKRETDYS...DGDPLDVHKRLP

BKb1 AQKRGETRALCLGVTMVVCAVITYIYLVTTVLPYQKSVWTQESKCHLI.....ET.NI
BKb2 ALKAGEDRAILLGLAMMVCSIMMYFLLGITLLRSYMQSVWTEESQCTLLNASIT.ETFNC
BKb3a SS.TGEDRAVMLGFAMMGFSVLMFFLLGTTILKPFMLSIQREESTCTAIHTDIMDDWLDC
BKb3b SS.TGEDRAVMLGFAMMGFSVLMFFLLGTTILKPFMLSIQREESTCTAIHTDIMDDWLDC
BKb3c SS.TGEDRAVMLGFAMMGFSVLMFFLLGTTILKPFMLSIQREESTCTAIHTDIMDDWLDC
BKb3d SS.TGEDRAVMLGFAMMGFSVLMFFLLGTTILKPFMLSIQREESTCTAIHTDIMDDWLDC

BKb1 RDQEELKGKKVPQYPCL..WVNVS.AAGRWA VLYHTEDTRDQNNQCSYIPGSV..DNYQT
BKb2 SFSCGPD CWKLSQYPCLQVYVNL TSS.GEKLLLYHTEETIKINQKCSYIPKCG..KNFEE
BKb3a AFTCGVHCHGQGKYPCLQVFNLS..HPGQKALLHYNEEAVQINPKCFYTPKCHQDRNDLL
BKb3b AFTCGVHCHGQGKYPCLQVFNLS..HPGQKALLHYNEEAVQINPKCFYTPKCHQDRNDLL
BKb3c AFTCGVHCHGQGKYPCLQVFNLS..HPGQKALLHYNEEAVQINPKCFYTPKCHQDRSDLL
BKb3d AFTCGVHCHGQGKYPCLQVFNLS..HPGQKALLHYNEEAVQINPKCFYTPKCHQDRSDLL

BKb1 ARADVEKVRAKFQEQQVFYCF SAPRGNETSVLFQRLYGPQALLFSLFWPTFLLTGGLLI
BKb2 SMSLVNVVMENFRKYQHFSCYSDPEGNQKSVILTKLYSSNVLFHSLFWPTCMMAGGVAIV
BKb3a NSALDIKEFFDHKNGTPFSCFYSPASQSEDVILIKKYDQMAIFHCLFWPSLTLLGGALIV
BKb3b NSALDIKEFFDHKNGTPFSCFYSPASQSEDVILIKKYDQMAIFHCLFWPSLTLLGGALIV
BKb3c NSALDIKEFFDHKNGTPFSCFYSPASQSEDVILIKKYDQMAIFHCLFWPSLTLLGGALIV
BKb3d NSALDIKEFFDHKNGTPFSCFYSPASQSEDVILIKKYDQMAIFHCLFWPSLTLLGGALIV

BKb1 AMVKSNOYLSILAAQK.....
BKb2 AMVKLTQYLSLLCERIQRINR.....
BKb3a GMVRLTQHLSLLCEKYSTVVRDEVGGKVPIEQHQFKLCIMRRSKGRAEKS
BKb3b GMVRLTQHLSLLCEKYSTVVRDEVGGKVPIEQHQFKLCIMRRSKGRAEKS
BKb3c GMVRLTQHLSLLCEKYSTVVRDEVGGKVPIEQHQFKLCIMRRSKGRAEKS
BKb3d GMVRLTQHLSLLCEKYSTVVRDEVGGKVPIEQHQFKLCIMRRSKGRAEKS

```





Figure 7

FIGURE 8A

```

1  aatcatgtga gttcatcatt ataaaacagc tgatttataa caactgattt taaagtgggt
61  tgtagcttg aagataaatg agatagggca tgattcaggt ttctgtgtac acactatatt
121 ctaatgaaag taattacaat tatcttctaa acagctgcag atattttttc tataaacaat
181 tttagatgtt acattgttaa atgttagctt tcaaatctct cacattttta ctcaatgaag
241 tccttttttag caaacttaca ggaatcattg tatcattcag ctattaaata aggaattgtt
301 ctaattcaca tcttaattaa gaactttact aggttatatc ttttgcaagt atgagaataa
361 tctagataga aggactagaa ttggattaaa ggtgatccag ataaaaatag taattctaatt
421 gaggaaattt tttaacattg aaaatagtac cctgtttatt ttttttaatg ttagagtcaa
481 accattgcac attactgtgt aaaaatacac aaagatggta acttgctgct aaagcccttt
541 ttacttttga atctttgtat ttttttcacc ttgttaattt taagtgtgct ttttatcact
601 catttgtttt ctatctttat ttctgtttgg gttaaagtta caaacagcaa gtttttgtgt
661 ttaaattcct gaaaatgttg cgatggaatt acagtaatgt gttacggctt gggccctctg
721 caggagtgtg tctgcagtc agcagtgaac acaaggttcg aacagcacac tatcagagct
781 aaacagatac tagtactgt gaaaaacata atggattcag taaacctggc agctgaagat
841 aaaaggatg agttcatttt gttgcaacat aaaaatgtta gttttttgt cagacttttg
901 ttaagatgag gttttaaatt gttgtactat aaatactttt tacactgaaa tcaacctgtg
961 gggtttaact gatgcctact tctatatttt taagtgcgta tttgaaagtg tcgatattta
1021 ttctgcaacc ttgcacacat catttatatc ccctggacct cactatcctt gtgtatgaat
1081 tgtgattttt aaaatgtttt gtggctttaa aatatctttg aagcagtgga agaaccattt
1141 cttcaagcag atttttatga agaagtccaa aacagggtgat cctctttctg cattcaacc
1201 acgaccattt ttttaatatg gtcatacat ggtattttga aatcactatt ttactagt
1261 agtgtttgtt taattgcctg tgaacaaaca aggcgatgct acctgccaac cagaaatttg
1321 atttggttaag cagtaccctt tatgtttatc tgtacggcat tattttatac tatttgttac
1381 cttgtaaagg tggcaatgaa tacatgaaaa taccagcatt atcgatacaa agtataattt
1441 tcacaaactt ggtatgcttg aaacacagat cgggatgtta ttagatgtga caaaagctaa
1501 tgatgttaag cctcgatagg accttcagtg gacttaagcc agttgaagga aacataaacg
1561 catcaaagct taagtccaag atgacataac attggaattt taaatgtatt tgctctttca
1621 gggacaaaga ccccttcatt tccctgtctg ttgctttata atggctactt cttggaatg
1681 aattaagttc aaatgcagag aattgccagc atataaaacg caagctcag aacctgagt
1741 ttgaactcag cccctgtgta cagcctccgt gtggcctctg ttttaattaga tcgtgctgct
1801 atagcagttc cttctagctc agttgctttg atgtagtacc caaatttttg cctaaaagt
1861 atttaattag taataatttt taaagatata ggatgttgaa caaagtatag cacaaagaag
1921 atgtgatttg aggattgtat aatcataatg tcctgggaac ttcttaagta aaagatcttc
1981 ttaaattgat ctcaggtctt tattttcttg tatcagccag agttgaacaa acttttgtt
2041 aaaaaaagag ccagatagta aatattctag gctttatggg ccataaagtc ctttggagct
2101 actcatttct acagttttag agcaaaagca gccatagata gataatagg aaatgaatgg
2161 gtgtggctgt gtttcaagaa aactttaaca aaagctggca gctggctaga tttggctgt
2221 aaggtgctat atgctgacct ctgcttatta actaaagtca gtattgcatt ctgttttgcc
2281 tgttcatatc tatgaagact tagaattcct attcatcctt tctgggattc aggtgccaca
2341 tgggcagaga aacgtggtt ctatcaaadc atctatataa aatactttat aagtgaata
2401 ttacttgaat cttttgagat gttacaagtt ttttttttcc ttgagtcag ctaatagatg
2461 gttcaatata tgaatgagtc ctttgctgaa atgcttttag acttcagact accctgaacg
2521 ttgattactc tttatactga aataggcatt attcagtgga agagagggaa gaccaaattg
2581 atagactgga ctttattcga aaccagatga accttttaac actggatggt aagaaaaaaa
2641 tcaaggaggt taccgaggag gtggcaaaca aagtgggtta cagtagcttc atgattaaaa
2701 taacctggat ggaaattatt tttgataatg ctaaaaatgt tattcctgtt acttcaaaag
2761 aatgttttca gatttttcaa ttttaatttt ttgaataatt gataatcttg ttttaattat
2821 ttaactccaa aattttcata ttttcaggtt tcatgtgcaa tgacagatga aatttgcga
2881 ctgtctgttt tggttgatga attttgttca gagtttcac ctaatccaga tgtattaaaa
2941 atatataaaa gtgtaagtta aagtatagat aaaattattc agagacaggt tcttattatt
3001 ctataccctc atttatttca tggtttggca tttcagtgta tcagtacaaa atgaaactgt

```

FIGURE 8B

```

3061 tagatctctt gtgccctctt gtaataaatg taaactgtct .tgtataaaaa gtaaatagaa
3121 aatttatact tagaatgaga taaagatcat tttaggacga ggcacagtgg ctcatgcctg
3181 taatctcaac actttgggag gccgaggtgg gaggatcagt tgggcccag agttcgagac
3241 cagcctgggc aacacagtga gaactcta ctacaaaaa cagttaggct ggttgcggtg
3301 gctccgcct ttaatccag cactttggga ggctgaggcg gacggatcac gaggtcagga
3361 gttcgagacc agcctgacca acatggtgaa accctgtcta ctaaaaaatac aaaaaaaaa
3421 ttagccggtg gcaggcgctt gtaataccag ctactctgga ggctgaggcg ggagaatcgc
3481 ttgaaaccgg aaggcagaag tggcagtga ccaagatcac accactgcac tccaacctgg
3541 gcaacaagag caaaactcta tctcaaaaa aataaaaaat agccaagcat ggcgacacgc
3601 ttctatagtc ccagttactc aggagtctga ggcaggagga tcgctgagc ctaggaggtc
3661 aaggctacag tgagtcaaga tcaaagcact ccagcctagg caacaagca agaccctgtc
3721 tcaaaaaaaa aaaaaaagtc aaattaaaaa gaccattttg gcatttactg aatattttat
3781 gtctttataa aaactacata ctttctggag aaaaaataat atggatattt accattgtta
3841 acaggaatta aataagcaca tagaggatgg tatgggaaga aatttggtcg atcgatgcac
3901 cgatgaagta aacgccttag tgcttcagac ccagcaagaa attattggta ctatttatgt
3961 ctacaaggtc atgtctggtt tgttttttca ttcagtactg gtgaagagct tattttcctt
4021 taggcattcc attgaaggta aaacatttac cattcttata ttaagtgttg taattttgtt
4081 ctttctagaa aatttgaagc cttacttccc agctggtata caggataaac tacatacact
4141 gatcccttgc aagaaatttg atctcagtta taatctaaat taccacaagt tatgttcaga
4201 ttttcaagag gatattgtat ttcccttttt cccttgggct ggtcttccct tgtacatcga
4261 tttttgggcc tagaaatgct caaagggtgc tcctaggatt atcagagcct atctttcagg
4321 tatgtatctt tgaatctacc aattaagact ctcttttatt attttgttta tgtggttttt
4381 ctataataaa actagcttta caaaatctgc acatttaaaa gctagtatct atctcttaga
4441 gccatttctg aggtagatat ttagtttgag catctccaat ctgaaaatcc ttaaagggtcc
4501 aaaaagaaa actttttgag tacggtgaca ccacagggtg aaaatccac accagacctc
4561 aatgaggtca tagtaaaaac gcaggcacac aacacacagt ttattcagtg tcctcaaggg
4621 aaaaaagact cagcctcttc tagctgcaat atatcttttc cacgtatgcc caaatccccc
4681 cacacaagca cacctatgaa gggactataa atggcacatg tgcaggtcca gtgcaccaac
4741 agcattttcc ccgatgccc ccacagggtg ccaagacctt tgtgcatcac tctggtttgg
4801 cttctttact tcctgtcttt tctctggtgt aaagatactg tttaaaaaaa attttttttt
4861 gagtggggat gcgtttgaga cagagacctg cgctgtggct caggctggag tgcagtgggtg
4921 caatcttggc tctactgcaac ctccacctcc cgggttcatt cgattctcct gcctcagcct
4981 cctgagcagc tgggattata cgcacccacc accacaccca gctaattttt gtatttttag
5041 tagacatggg tttcacctgt ttggccagac tggctctcga ctcttgacct caagtgatcc
5101 acctgcctca gcctcccaaa gtgttgggat tacaggcgtg agccacagca cccaggctgt
5161 ttaaaatggt taaaaggcct acagataccc ttgtgggtga cattgataag aaaaagaaga
5221 ggcattttatt tttatttata acacataaat ttaagctggt ggaaaaactg gacaataatg
5281 taagtgaaga acttcttaca gaagaatgat gttggaatga acaccatata tgacctaaaa
5341 gaaagaacaa ctttcctaaa gttgttcttt aggactaaac tgttgaagtt ccatgctaaa
5401 aatgatgaca gaagttaatg aaaaaaacaa aaacactgca gaaagttaaa aacgaagatc
5461 ttgatcatgc attgaaagag tggatccatc agcattgcag tgaacacgtg ccacttaagg
5521 acatgctgat catgaaataa acatctatta aaatgaactg aaaatcgaag ggaactgtga
5581 gtattcaata ggctggttag agaaatttaa ggtaagatag cattaacttt tctaaaggtt
5641 tgtggtgtta aagcatcttg atcacaaaac agcagagaaa ttcattgatg aatttgccaa
5701 gattgtctct gatgaaaatc tgactccaga acaagtctat aatgctgatg aaacatcact
5761 gttttgccat tattgaccca gaaagacact gactacagct gaggagacag cccctacaag
5821 aataaaggat gcaaagaaca gaataactgt gctgggatgt gctaatgcag caggcataaa
5881 tgtgaactta ctgtgatagg caaaagcttg catcctcact gttttcaagg aatgcatttt
5941 ttttactagt ccattattgt actaacaaaa aggcattggat ccctagggac atcttttctg
6001 attggtttca caaacatttt ataccagctt gtgtgcactt gcagggaagc taggccggat
6061 gtgactgca agattttgtt attccttgac aactattctg ctcatcgtct agctgaaatt
6121 cttattaaaa aaagtgtttc attatttcaa ccttgtgacc aggttattct acgatcaatg
6181 aagagtaaat acaaaaacct tttttgggca gcatgctagc agcagtgaac tgaggcctgg

```

FIGURE 8C

```

6241 gtgtggaagg ttttcagaag gagtttagaa tgaaagatgc catagatgct tttgccagca
6301 caaggagttt aggatgaagg atgccatata tgccagcact tggaagacag tgactcagtt
6361 gtgtatgcct gcagcctctg cctgtcacta cttttattga tgatgaggag caaattggtg
6421 actttgaagt atgtcaagt agctccttat gcaaaaaata atatcttcag agtccatccg
6481 taagctggaa gaaggatata tcaaagaagt gtttgacatt gataatgagg tttcagttgt
6541 tcattcatta actgatggca aaatagctaa aatgcttcta aatcaagggt attatgataa
6601 tggtgataat gaagatgatt tcattgacct cgcagaacaa ctgcctatag ggcagatggg
6661 gatgggctta ttgaagcact agagcagcat gtattcataa cagaacaaga aataggttta
6721 taaaatcaag gagagacttc aacaaaaacc attgtttaa aggcaagtag gtggtactgc
6781 aggaacatt taaaaggcc attcagcaca atgtcttatt atgccgagag gaccacttc
6841 ttggtccctc aactgcttct gatgtttctt ctacctaac aaagtactgt gtaccggaac
6901 tttttaatca aaacataaca ttgtaggtag agactgaaag cctgccattg ttgctgttta
6961 acagctgata caggtgttct ggtgatgcca ctgtgctgct tagtttgaat acgttatttt
7021 tctactgttat taatgggtgtg tcttatattt ttactattaa gttcctttgt gtgaatccgt
7081 gtaagaaaat gattgcttgt cagtagtatg taaattcaat caagaatgat ggtgatgcca
7141 aacaaccata gagtggtcac atgggtggct gacatagcaa cacctgtgtt ttctgataag
7201 tcagtgtaca caacctttgt ttcatgcaca aaattattta aatattggat aaaattacct
7261 tcaggctatg catataagg atataagca taaatgaatt ttgtgtttgg acttgggtcc
7321 catcctcaag atagctcatt atatctatga actattcaaa aatcctaaaa aatctgaaat
7381 ctgaaacact tctggtccca agcatttttg ataagggaca ctcaacctgt agtatgctta
7441 agggaaagct tatcctaaag ctttcgtcca atactgttgc aggtgggtgt acattcattt
7501 ttgaaacatt tctgttttct taaaagattt ggttttcaca ttcaaatata actagcataa
7561 atacaagcat ttttttaatt tttttttatt atacttttaa gttctagggg acatgtacac
7621 aacgtgcagg tttgttacat atgtatacat gtgccatgtt agtgtgctgc acccattaac
7681 tcatcattta cattaggtat atctccta atgctatccct ctccttctc ccatcccaca
7741 acaggccccg ggggtgtgatg ttccccctcc tgtgtccaag tgttctcatt gttcaattcc
7801 cacctatgag tgagaacatg cagtgttttg ttttttgtcc ttgcaatagt ttgctgagaa
7861 tgatggtttc cagcttcac gatgtcccta caaaggacat gaactcatca ttttttatgg
7921 ctgcatagta ttccatgggt tatatgtgcc acattttctt aatccagctc atcatgttg
7981 gacatttggg ttggttccaa gtcttttgcta ttatgagtag tgctgcagta aacatatgtg
8041 tgcattgtgc tttatagcac catgatttat attccttttg gtatataccc agtaatggga
8101 tggctgggtc agatagtatt tctagtctta gatccctgag gaatcaccac actgtcttcc
8161 acaatgggtg aactagttta cagtcccacc aacagtgtaa aagtgttctt atttctccac
8221 atcctctcca gcacctgttg tttcctgact ttgtaatgat tgccattcta actggtgtga
8281 gatgatattc cattgtggtt ttgatttgca tttctctgat ggccagtgat gatgaacatt
8341 tttcacgtg tctattggct gcataaatgt cttcttttga gaagtgtctg ttcatatcct
8401 tcgcccactt ttgatgggg ttgttttttt cttgtaaaatt tgtttgagtt ctctgtagat
8461 tctcgatatt acccctttgt cagatgagta gattgcaaaa attttctccc attctgtagg
8521 ttgcctgttc actctgatgg tagtttcttt tgctgtgcag aagctcttta gtttaatgag
8581 atcccatatt tcaatttttg cttttgttgc cattgccttt ggtgttttag acatgaagtc
8641 ctgcccattg cctgtgtcct gaatgatatt gcctaggttt tcttctaggg tttttatggg
8701 tttagggtcta acatttaagt ctttaatgca tcttgaatta atttttgtat aagggtgaag
8761 gaagggatcc agtttcagct ttgtacatat agctagccag ttttccagc accatttgtt
8821 aaatagggaa tcttttcccc atttcttgtt ttgtcagggt ttgtcaaaga tcagatagtt
8881 gtagatgtgt ggtattatct ctgagggctc tgttctgttc cattgggtcta tatctctgtt
8941 ttggtaccag taccatgctg ttttggttag tgtttggtag tgtagtatag tttgaagtca
9001 ggtagcgtga tgcctccagc tttgttcttt tggcttagga ttgacttggc aatgcgggct
9061 ctttttttgt tccatatgaa ctttaaagta gtttttttcc aattctgtga agaaagtcat
9121 tggtagcttg atggggatgg cattgaatct ataaattacc ttgggcagta tggccatttt
9181 catgatattg attcctccta tccatgagca tgggaatgtt ttccatttca cccattcaca
9241 attgcttcga agagagtaaa atacctagga atccaactta caaaggatgt gaaggacctc
9301 ttcaaggaga actacaaacc actgctcact gaaataaaaag agaatacaag caatttaaaa
9361 gaaaaacttt aaaaaattta tcaattcagt aatttttttag gaaatttgtg aaattgtaaa
9421 accatcacca caatttagtt ttagtttttg tcacctcaaa gctgttttac tcattttag
9481 tcaattatga tttccacctc cagcctcacc actaatctgc cttctgttta tggatttgcc
9541 ctttctggag gtttcctata aatgcaatca tatagaatgt ggccttatat gactggtttc
9601 tttcacttag tataatgttt tcaaggttta tcatcatggg atcagtgttg taggatgtat
9661 cagtacttca tttcttatta cggctgagga atattccatt atagatacaa cactatccat

```

FIGURE 8D

```

9721 tcaccagttg atggacatta gggtagtttg tagtttttgg ccattatgaa tactgtctata
9781 aatattcatg tacatgtttt ttatgtggac gtgcgttttc atttctcttg agtatatacg
9841 taggttctaa tgggtataga gtagtattcg ttgtaggttt gatttgcatt tccctaataga
9901 ccaatgattt taaacatctt ttttgtgtgc tcactagcca tttgtgtatc ttcttttgtg
9961 aaatgtctat tcaaagcttt tacctatttt taaattattt gggctctata gaggttgcagg
10021 atttctttgc atattctgga tcaaatcctt tctcacatat atgatttcca aatatgtttc
10081 tttatttttt gagacaagga cttgctatgc ccaagctgga gtatgggtgg gtgatcacag
10141 ttcactgccca cctctgcttc caaggagctg ggaccacagg catgtgccac cacacctagc
10201 taatttaaaa gaaatttttg tagacacaag gtctcactgt gctgtgctgg ctggtttcaa
10261 actcctggtc tcaagctatc ctctgccaat ggcctcccag agtgctggga ctacaggcat
10321 aagtcactgc acccagcccc aaatatattt ctcttgctg tggctgtgtg ttttgaagtt
10381 taagaggttt ttattttgaa gtcctttttt ttttttttt tttttttgag actgagtttc
10441 actctgttgt ccagactgga gtgcagtggt gcaatctcgg ctactgcaa gctctgcctc
10501 ctgggttcac gccattctcc tgccctcagcc tcccaagtag ttgggactac aggcattccac
10561 caccctgctg ggctaatttt ttttttttat gtatttttat tagagacagg gtttctactgt
10621 gttagccagg atggctcga tctcctgacc tcgagatcca tccacctcgg cctcccaaa
10681 tgctgggatt acaggcgtga gccaccgcgc ccggccgaag tcctttttt tttttttagt
10741 ttatcatttt ttgttgttgt ttgtttcaag gattgtgctt ttgttgttag tttttttaa
10801 aacacacttt gtccctcttg agagttgagg gcctactgtt acatttagac cccattgtat
10861 attatgaaaa tttattgttc atgatcatat tgggtccat tgtagataag aatttccaaa
10921 ttatttttat tttctacata tatttgtggt gtactcatca tttagggtaa tttttccctc
10981 atagatgtgc tactataatt ttagagtttt ataaagataa ctttatagtt gtttaagtgt
11041 aatctttttt ctttctctt tttttttggc agctccctag atctttagct tctactccca
11101 ctgctcctac cactccagca acgccagata atgcatcaca ggaagaactc atgattacat
11161 tagtaacagg attggcgtcc gttacatcta gaacttctat gggcatcatt attgttgagg
11221 gagtggtaag aaacattact tttagtataa ttaaaatcga aatgtttgca aggcctgcctg
11281 ttagctcaca aaagaaaagg tatacagtat ttacttctat ttataggatc tgtaaaatag
11341 tatgaaaatt tgcaggttaa aagtagtgaa ttaaatatca caagtttcat cttaaatttt
11401 taaaaaacat acatcttttag gaatgaacta tcaccaccta gctggctctt aatacttctc
11461 aatcattttc acatcacagc acacaaaaga atatttgtac gttgaggtaa taagaaaacc
11521 ccagggtgcc tatctagcct tctctgaata tcaaggggat ctgtcttaag tattaatgtt
11581 tttttttagt agtagttctt atttgtttct ttaaaaaatg tatacatta acagcatgga
11641 tatgcttgtt ttgggttttg aaaggctcga cctacaatat ttaacattgt tttatttttg
11701 agatacattg aatcaggata agtctggcag cttataaagc tctccctcat ttgtgtgctg
11761 atccctttta agcctgattg atttctggta acttcagggt ttctcatgga gaaggaatat
11821 acatttttag aaaatgtatc taactcaaag atccataggg aactaaaatg cttttaaatg
11881 tactctccaa agtgggtgtt tctctctca gactaagcta tgactttatc ttacagattt
11941 ggaaaactat aggtgggaaa ctccatctcg tttcattaac tatgtatgga gtttgtatc
12001 tttatgaaag actgagctgg accaccccat gccaaaggag cgagccttta aacagcagtt
12061 tgtaaaactat gcaactgaaa aactgaggat gattgttagc tccacgagtg caaactgcag
12121 tcaccaagta aaacagtaag ttggaagggt catctttcct ttaaaaaaaa gttactgaaa
12181 tatgacatac atgcagaaaa agcacaaaat aagtgtattg ctcaagaat tatcacaaaa
12241 tgaacatgtt tcgtgatggc cataaatgag aaaaaataga acaatactaa acctcactgc
12301 tgccctttct cctcctaatt cactattctt ctttccattc tcctgataga ttaggtttga
12361 acattagaaa attggtagat aggaactctc aagaactctg gaggggttta aaaagatagt
12421 tcttaatttt ttttctttt tttcagagat agggctctctg tcgcccaggc tggagggcag
12481 tggcacaatc tggctcactg cagcctcgaa tcttgggctc aagtgatctt actgcctcag
12541 cctcccacgt agctgggacc acagggtgtg gccaccacac caggataatt ttttaatttt
12601 tttttttct ttgagacagg gactcaatat gttgccggg ttggtcttga ccacctgggc
12661 tcaagtaatc ctccctcctc aagcctcctg agtagctgag attataggca tgagccacca
12721 tgcccaactc aaaagatctt cagcagacct attctaaatt tatgtacctg gctgggcaag
12781 gtggctcacg cctataatcc cagcacattg ggaggctgag gcaggcgat cacttgagg
12841 cgggagttcg agaacagcct ggccaacatg gtgaaacccc atctctacta aaaacacaaa
12901 aattagccgg gcatggtagc acatgcctgt aatctcagct agttgggagg ctgaggcaca
12961 agaatcgctt gaccctggaa ggcagaggtt gtagtgagcc gagatcacat cactgaactc
13021 cagcctgggc gacagagtga gactctgtca cacatacaaa aaaaattaag cactggatat
13081 agatttattt ttctattctt tgtctttttc tccttagaaa gtgaacaga aaaaaacaaa
13141 aataaaataa cttctaattg attaagaatt caggttattt gtgttcttat taataggggt

```

FIGURE 8E

```

13201 tattctataa catttaggaa tgcataaaaa ttcattgatca gatatacactt gccagaagaatg
13261 ggggcttcat cagatccgga atagaattta tctaaaagtg atcaagacat gcagacttat
13321 aaaaagctat gaacatcctg tctgtataac aacttggcca gcaacattcc tggcgcaaag
13381 ggctaaggct ccttcaagcc ttgagaataa gacacttaaa agaataagcc caagctcctc
13441 ctgagcgagg aggccgaata ttgtcagtag aagcatggac ttttggatgt gatctgttct
13501 ggagccccgg acctagccct tgttactttg tgatttttgc acaagttcct cggactctct
13561 ggtcttctgt gtctcatctt gctgaatgct caaaaagttc ctacctctc aagttctgtg
13621 tctgaaggac attatgttct catagcactg agcacaatcc ctggcacatg gttactcagg
13681 gcacccaagt tatcattatg tgtctagggg aagttgggtt gggcatgcag ttgttgaatt
13741 ctcttcttct tgggtgagcg ctgcctctca gcagctgatg ggggaatcct tgcattattg
13801 tcaactccagg agagaagata cctgcttccct gcaagcaaac ttacggtttc atacacttta
13861 ttgatctca aaggcagatc tttttttggt ttgttttgct ttcttgagat ggagtttcgc
13921 tctcgttgcc caggctggag tgcaatggca cgactctgat gtggctcact gcaatctctg
13981 cctcctgggt tcaagtgtt ctccctgcgt agcctcccaa gtatctggga tgacaggcat
14041 gcgccactat ggccagctaa ttttgtagtt ttagtagaga tgggtttcac catgttggta
14101 aggtcgttct cgaactcctg acttcaggag atctacctgc gtcagcctcc taaagtgtg
14161 ggattacggt ggtgagccac cacaccggc ttaaaggcag atcttaaaag cacattaaat
14221 catctgctct aactcccaca ttgcacagag aaataagctg agttccaagg aggcagcata
14281 acctgagact agaaatggtg cttagggttc taagcccagg cctccaagggt ctatttactg
14341 tctaatgtga gctgatgtct ccaaaagtat aaatgtaggg tcacctgtgt taggatcata
14401 tgaagggtt tttaaaatgc aagttcttgg gtcccttacc ctaattgttt tatcagatc
14461 tctagagatg ggacctggga atctgtatct aacagaggat cacacctgag ttcaagaacc
14521 actcatgtag tagaacaatt acctcaactt aaaatatgaa atgtatctgt agcaagtgcc
14581 acctggtaaa gacttgatca cagtggattt caaacaagac aaagtattga gggctgttga
14641 actgtcaaaag aatttcagct attatttcta ttagtttctg cctcactatc catcgagttg
14701 tttgtatgcc aggataggcc aagttctctg gctcttgtga gcttgtgtaa gtcagtgtag
14761 ctctctgctc ttgaaaaagc ccaaaaagtg aattaacatt tgtatagact tagaactgta
14821 actagctcat aaatagtatg cactagtatt atcactcaga gcagaaaaag catctgcaca
14881 gaggtgacgc tggtttctct gatgtgagcc tagttcaggc agtcagggtc ccatttgatt
14941 agcaagatgg ctggagataa taatcagggt aaaggaaagg aaagatccca tcgaaggctc
15001 cagaagtttg tggcacagat catacttccct tgttgtttgt tttctgcaa agagaatgta
15061 agatcagatg tgactttact tttgacagtt tgaattcttg ctacatcag caaaaattct
15121 ccataattgga ataactgtcc agttaggggt ttacttattc tcctacgaac aaatatagat
15181 agtcacgcaa agaacatagg ctgttgtgta aattttaggt tgttgtcaat gatttgtgca
15241 tctctaaatt ggaaaacaca gacatagttt tcatgaacat ggagaatttc agtcacaaa
15301 taattcttag ccataatagg tattgtatat ttaattgaga gaatgtgaaa aacaatgagg
15361 aagtagtttc tccagtatgg tgaaaggcaa agagggttct ttttttccc ctaagtaaag
15421 catctactaa atgcaaaaaga aatgattgtg gactctggaa tctggaatcc acgatgctag
15481 cactttgcag taatagcctc tttcatatat agatctcaca acagtttgta gacactaagt
15541 ttttccatgc tctgcaact cacactgagt ttttatttac tcctttttta ctctggaaaa
15601 gcagggcagg aagtttttaa agggttcctt gcagttacaa agctagaatt tgaaccaggt
15661 gtcaaagcct ctgggtcaga ttgggagggg gccacggtg tgaacttgac agttaattca
15721 tggttgctgt tttatggagc aggaagtgtc ctttagtgac tcagaggaag gaataagctg
15781 agggagggtc caggagacca gaggtacagt gcctggcatc tcttatcact gcattctaag
15841 tggcttagcc ttgctgtgtt cctgaatgca cacattggga ttgcaactca caccgaccac
15901 tttaggttct tcttgagaga taggctgtga gtccctgaatg gagtgtgtgt gtgtaggggg
15961 cttggcttct aggccaaagag tggagtgtgt gtggtacttg gtggacagag aggccagttg
16021 gttatcaagt gaaggggctg tagggaaaaa gttcagaggg caaaaagcta tgttgcctct
16081 tctcacttat tgtccatgag tattttcttt taaatggaac cggaatcata agataagaat
16141 gaccaggagg ctctggtaaa ggcagggtgca agttttgtgt gacaagatcc tcatggacta
16201 aaaaatgatc atatttctga ggttgaatca aatacataaa agcagcagca aatcacacac
16261 ttcttcgaag agagttacca agttgcaggg gaaatttatt agcacaata agtctgaaaa
16321 ggaaaagcta tccatatata caaggttaag ttggaagaag gaggcagaaa gaagaaattc
16381 cagttgttat cattattttt tattccaaat tgcattctgt cactaaaaata cttcggactg
16441 ggtgcagtgg cttacaccta taatcccagc actttgggag actaggcggg cagatcactt
16501 gaggtcaggc attcgagacc agcctggtca acatgatgaa accctgtctt tactaaaaat
16561 acaaaaatta gccagtcgtg gtggcacatg cctgtaatcc cagctactcg ggaggctgag
16621 gcaggagaat cgcttgaacc tgggaggttg aggttgcagt gagccaagat tgagccagtg

```

FIGURE 8F

```

16681 cactccggcc tgggtgacag agggagactc catctcaaaa aaacccaaac aaatatttta
16741 taaaatagaa atagaaaaat aatacaaatg aaaagtctat taagtatata tttttattaa
16801 gcaatattaa aatgactgca agtaagagtt tatatagcta tatgtacatg gatatttata
16861 gatgagatta cgcttacatt cttgcccaca ccactcttgg aaatgttaag ataatatcgc
16921 cttgactgaa aaacatacag caaacatggt ctctttggca ttctgtcatc cacatctaca
16981 ggtgcctgta gcaatgtgtg gtactataat ataatggtaa ttgatgttcc taatttggga
17041 gtgtggaaag atcccaaat gtcttttaag tcatcagaga aagataaaat aatatttgat
17101 acagcttttc ttaaaatttg agataatttt aatggcgagt tattttatgg ccctttgtat
17161 cttgaaaaat tgggaaatca catatggttt aaaagcgaat tatcttaatt ggaacatgcc
17221 attaaactaga aaaccatta tttcagcttg cactctaaca gacaatacgt ggaaaaggaa
17281 acgcggccag ggcaaaacat ttctctctct tataaacctt gaactgagta cgtccctcac
17341 caattataga ggggtcccctt gggcctcaga actttccaca agcgttgagg tctctatggc
17401 gatgctcccg gctgccgagg cggaaacaca ggtgatgagg tggcggcaag cacagtgcaa
17461 agagagagaa gcagcttcgg ctgcagcaaa ccacgcaggt ccttcttgat cctctagaac
17521 tgaccgctcc gccttgccag gagtctgcag aaccacgtgg ctagcctgcc tgaagtcttc
17581 acctctccag gaaggcgggg ggcttcta at ggctgcagct gcgctggggg ctgggggctc
17641 ccgctgggac tccacttccg tggatgtcta agcttcacct ttcttgcgcc cgcaggggca
17701 tgactcaggt gaaagggagc cattttctca gaccctggc ctcatgcagc ccttcagcat
17761 ccccgtagaa atcacacttc agggcagccg gaggcgccag gggaggttaag tcacttccgg
17821 aagctctgcc ggtagtggga atctggctga acaagcagtt gcaagaagag gggacatctc
17881 gagcttgggg agtgagtgtt tcctttttct ctgaggatgc ccacttgcca tgcctccag
17941 ggtaccagc aggttcccc agtagcactc acatcacggg gctgcagcct ttctgttg
18001 ctctatcttc taggttgcca gttcttgagg actggagacc ttttaaactc acctgtagct
18061 cccagcact gatcatagcc cagcccatag ttggtgctca agaatactt gttggttgaa
18121 tgaggaatga agaaattaga ccagcatttg gtcccatttg tgaagccctg gagtcacagc
18181 ccttggtatc aaaccagct caccacttaa tcagccatat gactgggcag gtcccatttg
18241 tgaagccctg gagtacagc ccttggtatc aaaccagct caccacttaa tcagccatat
18301 gactgggcaa gtcacttaac ttctctgttt gcctcatctc cttatctgtg aaatgcagat
18361 agtaagagcc cctgcgtgtc tcagcacagt ataccacatc ctctctaaac tatctgtt
18421 taccgggtat acgctctaca cctccctaaa ctctagcctt ttagctattg ttattacca
18481 ccctactctt ctctttaaag ggggaagtga gagattattt tcaggaccctg tttctctccc
18541 agggaaatga aaagcaaaga agctgaaagc ctctagttag ctgcaccctt tttgcctgcc
18601 ctagggttgg ggcattgggg agtcaatgtg ggaacaaaaa tggagagaag agtgatgcct
18661 gagggtgttg gacaaatggg atactaaaac cttttgtgct aggcgcggtg gctcacacct
18721 gtaatcccag cactttggga ggccgaggtg agtggatcac ctgaggtcag gagttcgaga
18781 gcagcctggc caacatgtgt aaacccgctc tctactaaaa atacaaaaat tagctgggca
18841 tgggtgtggc gcgcctgtaa tcccagctac tcaggaggct gaggctggag aattgcttga
18901 acctgggagg aggaggttgc agtgagcaga gattgtgcca ttgactcca gcctgggcca
18961 caagagcaa actctgtctc aaacaaagaa aaacctgttg ggaggggtac atttcagaac
19021 caggttact cactatctgg aaatatgcat gatttattat tggctctagt ggagttggga
19081 gctgagaatt ggaaaacatt aaagatggta atgggtcatc tggtactcac ttccattatc
19141 acttaactgc actcaggggt atttgagagg gagcatgagg agagtgaat acaggagttt
19201 ggataaatg ggggttcagg gaagaaggac cagacagact acagagaaa gaaaggtgtt
19261 cttctcgcta gacatgaacc aatttttttg tgaacagaca attaaaatga attactttat
19321 ggcaaaagat caaatgacaa acatgcaagc aaacaagttt tagtgtccca tacgtcacac
19381 aattaactag atataaaggc agttgtgttg tcatcaaagc aaactactgt atccattttt
19441 catttctgaa atgcacaact gaattattgc tatttctctt tgctgaactt gatgaactat
19501 gttgacttaa ccttattttg tgtttcaaaa taagttgtta aataatgttg taattaaaaa
19561 atagaagagt aaaaataatt accaagggtc tcctctgtaa ccaaacgaa ttacagggaa
19621 atattaatat agtgacttc atgttttagc attcatttca catacaccg tgaatgaat
19681 gtcagatgaa ttatgaagtt ttgatgagcc acagcatgtt tcttcttgca
19741 aaagtttcag tgctggaggg agaggctgct ggcttctgtg gggatcacac ccaggtgagt
19801 gtgttcaggc tgtttgtaat tgagtttgcc tcaggctaag ccagaagctg cctgtagcca
19861 tgtgtcttac ttgggctggg tgggaaagtc agtctatct gcagtgaaga gaatagaagg
19921 tgggtgatat tgccttgatt atgaataaaa cagctcaagg taatacactg gttagaagcg
19981 gacatgtatt actggcagaa aaaggaatca atatgctttt tatccatctt cctgactaga
20041 aagcaacta gatcatactt aagtgttttg aggtccttgg atgaaagatg cttgtaaata
20101 caacaaagtt aattacaagg ctgtttatgg tctgagaaaa ctggaacaa cctaataata

```

FIGURE 8G

```

20161 ttcaataata aggaaatggt taaggaaata tggcatatct aattgatgga acattatgta
20221 gccaatagga ttacaaagaa ttgttaatga catgggaaag tgcttattat gttaagtga
20281 aaagataaga ttaacaaaaa aatctcaaat catatcta atgtatctcat tctgtttaa
20341 acaatatagg ccagggtcgg tggattatgc ctataatctc agcactttgg gaggccgagg
20401 cagggtggatc acctgaggtc aagatttcga gactagcctg gccaacatgg tgaaccccg
20461 tttctactaa aaacaaaaaa attagctggg cgtggtagcc ggccgctata atcccagcta
20521 ctggggaggc tgaggcagga gaattgcttg aaccgggaa gcagaggttg tagtgagctg
20581 agatcatgcc actgcactgc agcctggatg acagagtgg actccatctc aaaaaaaaaa
20641 aaaaaaaaaa aaagaaaaga aaaagaaaaa caaacagaa acaaaaaaca aaaaacaaaa
20701 aacctaatat agagcaggag gggattaacc cagcaacca agtgacaat cattaccttt
20761 aagtgtgtg ggtgagtttt tgttctcttc tgtacatttt ttttgtatt tttcaagttt
20821 catacaatga gcatataaaa atatattact tcatgaatc atttgacatt tgttgaggaa
20881 ttctttgtgg ctaagtttgt agtcaggctc tgagaggtga caacgtgctg gcagcccttg
20941 cagccctcgc tcgctctccg cgctcctcgc gccttggcgc ccactctggc cgcgcttag
21001 gagcccttt ctgggctggc caaggctaga gccggctccc tcagcttgca gggaggtggg
21061 gaggagagg cgcgggcggg aaccggggt gtgcgaggag cttgcgggcc agtccgagtt
21121 ccagggtggc gtggactcag cgggccagca ctccgagtg ccgaccggcc tacaagccac
21181 ggacagtgg gggcttagca cctgggcccag cagctgctgt gctcaatttc tcacagggcc
21241 ttaggtgcct ccccgcgggg cagggtcttg gacctgcagc ccgccatacc tgagcctccc
21301 cccgctcctg tgggctcctg tgccgcccga gcctccctga tgagcgctgc cccctgctcc
21361 acggcaccca gtcccatcca cactcaagg tctgaggagt gcgggcacac gcacaggact
21421 ggaggcagc tccacctgtg gcccgggtgc gggatccact gggtaagcc agctgggctc
21481 ctgagctcgg tggggacttg gagaacgttt atgttttagc aagaaatctg aaatacacca
21541 attggtactg tgtatctagc tcaaggttta taaacacacc aatcagcacc ctgtatctag
21601 ctgagggttt gtgaatgcac caatcgacac tgtatctagc tactctggtg gggacttggg
21661 aaacgtttgt gtccacactc tgtatctagc taatctagtg gggatgtgga gaacctttgt
21721 gtctagctca gggattgtaa acgcaccaat cagcacctg tcaaatggg ccaattagct
21781 ctctgtaaaa tggaccaatc ggctctctgt aaaatggacc aatcagcagg atgtgggtgg
21841 ggccagataa gagaataaaa gtaggctgcc ccagccagca gtggcaacct gctcaggtcc
21901 ttttcacgc tgtggaggat ttgtctcttt gctctttgca ataaatcttg gtactgctcg
21961 ttctttgggt ccacgctgcc tttatgagct gtaacactca ctgcgaagg ctgcagcttc
22021 actcctgaag ccaggagac cacgaacca tcgggaggaa tgaacaactc cagaggcgcc
22081 gccttaagag ctataacact cactgcgaag gtccgcggtc tcattcttga agtcggtgag
22141 accaagaacc caccaattcc ggacacagtt tcataaatgt tccatacatg cttgagaata
22201 atatatattc tgtagaagtg agtattctat atttatcatt tagataaaa ttgttaattg
22261 cttacttaa atctattacc ctactggttt gttcagggtc agctatctaa attactggg
22321 gagtgtataa aaatacatca taatctgata gtggattttg tctattcttt cttgtagttt
22381 aatcagtga acaatgctat cagggtaccta caaattagca ttgttacatt ttcggtgtga
22441 attgagcctt taatcagtgt taaaacactt atttttaaat ttctaataaa gctttttaat
22501 ttttaacatt catgtttgct ttttgttaca ttttgccat aaatttcacc cttttgtat
22561 ctgtgtttta gatgtatctt ttgtaaaaac atgcatatag atagtctctc atttacagt
22621 ggattacatc ctgacaaatc catcataagt ggaaaatact gtaagtggaa aacttcagag
22681 tattagctat ttaccctcat gattgtgtgg cagactggga actatggctg gctgccactg
22741 ccagcatct caagagagta gggactgca tatcgctagc ccaggaaata atgcaattc
22801 aaaatttgag gtacagtttc tactgaattc atattgcttt cacaccatag taatgtaaaa
22861 aaattgcaag tgtggggcgg ggccggccat gggggcggcg gaggcgcccg agcccgctt
22921 ccccgcgcc attccacccc cggccaggct cagcccgcg cgctccccg cggctgggt cttggccccg
22981 ccggtgcgc ccgagcccag tcccgcgagc cgctccccg cggctgggt cttggccccg
23041 gaagcgcgag cgttcacttc gcggcgagtg gctccgtctc cgggacaga gcgcgcgcc
23101 cctggcccgg cccgcggggg ggggctccgg cacggtcccc gagcgtttc cgcgggtgg
23161 agcgggccga gccagcagg ttgaccagcc ccgcccgcac gcagcgccg gagatctact
23221 gtttgagcgc ggaagcgag aggtggcggt aggccgggt cgccgcaaaa cgggagccc
23281 gcgcggaggc tcgtgagatc cccatgaagg agctggagcg gcagcagaag gaggtagaag
23341 agagaccaga aaaatatatt actgagaagg ggtctcgtaa catgctgggc cgtctgcag
23401 ccacgctggc ctttctgggt gggacttcct ctgagagagg cagcgagac acctctatat
23461 ccatcgacac cgaggcgtcc atcagggaaa tcaaggactc tctagcagaa gttgaagaga
23521 aatgtaagaa ggctatgttt tccaatgctc agttagacaa tgaacacaca aacttcattt
23581 accaagttga caccctgaaa gatagtgtgc tggagattga agaacagctg gctgaatata

```


FIGURE 8H

```

23641 ggcggcagta cgaagagaaa aacaaataat ttgaaaggga aaaacacgcc cacagtatac
23701 tgcagtttca gtttgctgaa gtcaaggagg ccctgaagca aacagaggaa atgctcgaga
23761 aacatggaat aatcctaaat tcagaaatag ttaccaatgg agagacttcc gacactctca
23821 gtaatgtttg ataccaagat cctaccaaga tgacgaaaaga agagttaa at gccctcaagt
23881 cgacagggga tgggacccta ggaaagccag tgagggtggag gtgaagaatg aaatcggtggc
23941 gaatgtgggg aaaagagaaa tcttgacaaa tactgagaaa gaacaacaca cagaggacac
24001 agtgaaggat tgtgtggaca tagaggtatt cactgctggg gagaataccg aggaccagaa
24061 atcctctgaa gacactgccc cattcctagg aaccttagca ggtgctacct atgaggaaca
24121 ggttcaaagc caaattcttg agagcgcttc tctccctgaa aacacagcac aggttgagtc
24181 aaatgaggtc atgggtgcac cagatgacag gaccagaact ccccttgagc catccaactg
24241 ttggagtgc ttagatgggt ggagccacac agagaatgtg ggagaggcag cggtgactca
24301 ggttgagag caggcagaca cagtggcctc atgtccttta gggcatagtg atgacacagt
24361 ttatcatgat gacagatgta tggtagaggt cccccaacag ttagagacaa gcatagggca
24421 tagtttagag aaagaattca ccaaccagga agcagctgag cccaaggagg ttccagtgc
24481 gagtacagaa gcaggtaggg atcacaacga agaagagggt gaagaaaaag gattaaggga
24541 tgagaaacca atcaagacag aagttcctgg ttctccagca ggaactgaga gcaaggggtca
24601 ggaggcgaca ggtccaagta cagtagacac tcaaagttaa ccctcagata tgaagagcc
24661 agatgaagaa aagaatgacc aacagggaga ggcattggac tcattgcaga agagaaagaa
24721 caagaaaaag aaaaagaaga ggaaaaaatc cccagtacc atagaaacc ttaaagatgt
24781 ttaaaaagag ttaacttatc agaacacaga ttaagttaa ataaaggaag aagagcaggt
24841 aaagtctact gacagaaagt cagcagtggg agcccaaac gaggtgactg aaaatccaaa
24901 acagaaaatt gcagcagaaa gcagtgaata tgttgattgt ccagagaatc ctaaaatgaa
24961 gttggatgga aaacttgacc aagaaggcaa tgatgtaaaa acagcagctg aggaagtact
25021 agctggtaga gacacattag attttgagga tgtcacagtt caatcatcag gcccgagggc
25081 tgggtggtgaa gaattagatg aaggtgttgc aaaagataat gctaaaatag ctggtgccac
25141 ttaagcaat cctgaagaac cagagagcga agatgcagat cactgcaccg taccaaaaa
25201 tgaaagtccc tcacaggaca ttagtgatgc ctgtgaagca gaaagtacag agaggtgtgg
25261 gatgtcagaa catccaagtc agaccatcag gaaagcttta gacagcaata gcctaaaaa
25321 ccatgacttg ttggcaccag gaggagagcc gggggacttc aatccagaaa ctagagaaga
25381 ttgacagga ggaacgaga agggcaaaag caaagaagac cgtaccatgt cctaaagtga
25441 ggcaggcggc aggcgtggtg cacaggaagt ctgagtgtga ggggctctt tctctccact
25501 gccaatgtaa gtagaatgtt ctaaattcat agagatgcac tgtatgccaa tcaccaggtg
25561 atctactgct ttaagttata gactgttact tgtagatttc catgtaatca ttgaggttat
25621 caccagatt agaaagacat atttgttatc agtgtacatt ctaattgaga gcataatatc
25681 agtagtatca aacaataatg tctactgttt atagtccact taataaaaaa agaagcattt
25741 accatttgcc ttaggtgat aggaattgta atattcttga ccaaatatat cagcatctaa
25801 ttgaaatgac caaatagcat tcttagactt ctgtattatg aatataattg atatttaaat
25861 taatgtcttg ttcatatatg tgtactttca tatttgattt taaaatatac attataacct
25921 gtatggtatt ttatttaaag gagataaacc gccaaatagc aaatagggtca ctgaaaagat
25981 ttgcacctta gaacaataat cattttaagg ataacaagta aaggtctgaa agcatgaggg
26041 gctttatttg ccttcacctc atataagctt ttgattttga accaatgctt ttggtctca
26101 ttgttgatga tacttgaatt tactttgtag gagattttaa cttcatgctg atgatgtatc
26161 aaattcattt tatacaaagt ttaaagattt tttctggaag tgatacatgt caaattacat
26221 ttcctactgc agtatttgag caggacagct cattttttaa atgttttttg ccgggtgtgg
26281 tggctcacgc ctgtaatctc agcacgttgg gaggccaagg cgggtggatc acctgaggtc
26341 agaagttcaa ggccagcctg gccaacatgg tgaaaccctg tctctacgaa aaatacaaaa
26401 aattggccag gcgtggtggg gggcgctgt aatcccagcc actccggagg ctgaggcagg
26461 agaatcgctt gaacctgcga ggcggagatt gcagtcagcc aagatcaagc cattgtactc
26521 cagcctggac aacgagcgaa actctgtcta aaaaacacac acacacacac acacacacac
26581 aaaacaatgt tttcatgcct gtaaccctag cacattggga agccaagttt ggaggatcgc
26641 ttgaggccag gagttcaagg ctgcagttag ctatgattgc accactatac ttagcctgg
26701 gagacagagt gagaccctgt ctctaaaaaa aagaaaaagt ttttgaacct taaaattact
26761 tctctttgtt tgaatttcta atcatcattc aaaagaacag ttaaaaaagg ttacttgttc
26821 ttgtgcaact acaaattaga ctggagtagg atatttttaa gagctgaatc acttttggta
26881 ttttgttata aatgttttca tttgttatgt cccagtatat tcttattgga aaattcttgt
26941 tttgatctgc ctgaagaaaa tatctgtttt ctatataaag aaacatttaa aaataattgt
27001 aaagttagat ttaattgtaa aatataaaat cacaaaggaa tgtaccttat gaatgttgac
27061 attttatgaa attatgtaga ttcattattac tgttacaaga tagaattgaa tgcaaaaaga

```

FIGURE 8I

```

27121 ccaaatcctc attaaaaattt gaggaanaaca taagtgttat tatgtaattg aaataaaaaac
27181 attttatagt tgtaaaaaaa attgcaagtg gaaccatctt aagttggggg acatctatat
27241 gtatttaaat ctagtctgac aatctttata tttgaaaaac agttttttta gagatagggt
27301 ctcacctatc actgaggctg gagtgcagtg gcacaatcaa gcttattgca gcctcaaaac
27361 cctgggctca agcaatcctc ctggctcagc ctctgagta gctaggacta taggtgtgcc
27421 actacaactg gatgggtgtt taatttttat tgtgtagaga caggggtctg ctatgttgcc
27481 caggctggtc tcgaactcct ggggttcaagt gattctccca cgttgggttc ccaaagtgtt
27541 gggactatag gcatgagcac cacagcctgc cctactctct atcttttaac tggatcattt
27601 actccattta gttttattgt aattactgat atactgatgc aataacatta ttctatcatg
27661 ttattctgtg ctatttgtcc tgactnttcc atgagttttt cctcatctt tattgccttt
27721 tttggattga tttttccct ttcattctt tgtttctcta ctagtttggg atttctggag
27781 tatcacctaa aagagtagag aaagggtgata tttctattta gcatgcatat ttgaactttt
27841 caacatgaaa ttaatgtctt tatttttccc catggaaaaa ttatcactgt tactccctct
27901 aaattatatt ctcttgtcat gtatttatct tttcttttaa cccacacaaga cataatagta
27961 ataagataat tactattatt attattttgt agaattaata tctttttttt tctttgagat
28021 ggagtccttc tctgttgccc aggcctggagt gcagtgggtg gatcttggct tactgcaacc
28081 tctgcctcct gggtttaggt gattctcctg tttcagcctt ccgagtagct gggatccacg
28141 gtgtgtgcca ccacgtccag ctaatttttg tattttttgt agagatgggg tttcgccatg
28201 ttgggcagtc tggctcggaa ctctgacgt cagggtgatct gcccacctgg gcctcccaaa
28261 gtgttgggat tacaggcgtg agccactgtg tccagcctag agttaatatc tttacttacc
28321 tgtttacttc tgtatttgc tttcatttct ggtaaaatat cagctgttat ctggattttt
28381 ttcgttctgc cttgagtaaa tacttgacct ttttcttca gtgagggctt atcaaggaca
28441 aactcagttt tgtgtttgaa atgtctttgt ctgaaatgtc tttattcatt gagaggattt
28501 ttttgtagac atagaattct gtggttactt tttctcagta tattgtaaat attcttgctt
28561 tatggcttct agtcttgtta cagagaagta agctctcagc caaattgtca ttactttgaa
28621 gttaattgtc tttttctttg gcgactgtta agatttcttc ttgtctttgt agctgtgcaa
28681 tatatctgtg atgtgtttta gtgtggttcc tctttatttt atcccatggg cttctggagt
28741 ctgggaactg gtcttcaatc agttctagaa tttgactatc ttttaaaat attgtctctg
28801 acttatgtct cttttcttct ggaattctga atagatatta tgttacacta tttaatctat
28861 ctttcatgtc tctgaacctc tcttaataac tttccattta aaaatctctc tgtattatac
28921 tctggctatt tttgcagatc cagctctgtg ttcactaatt atcttctcag atgtattctaa
28981 ttgagtgtta tgtctgttca ttaaattttt actttcaatt attttatttt tgatttctat
29041 aaattctttt ttaaatatta gctctacatt ttagaattcc ttgatttctg acattttgga
29101 tcttcccttt atttctttta atagtgtaca gatgtgtatt ttatattcta tgcaaatatc
29161 tgaaattttt cgtggatctc attttgtggt ctattttttt tgcactcaaa cgccttgatt
29221 tctgtgtgtt ttatgactt ttattttatt ttttattatg tatttattta ttcatTTTTt
29281 tttgagatag agtcttgctc tgttgcccag actgaagtgc agtggcacga tctcagctca
29341 ctgcaacatc caccttctgg gttcaagtga ttgtcccacc tcagcctcgg gagtagctgg
29401 gattacaggt gcctgccact acacctggct aatttttgta tttttaatag agacagggct
29461 tagccatgtt ggccgggctc gtctcaaact cctgacctca ggtgatccac ctgccttagc
29521 ctctgaaagt gctgggatta tagccatgaa ctacctgcc tggctgttgt gtgttttatg
29581 acttctaaat gagtgcatat actccttgga catttatata cgtcttcttt ttttatttta
29641 ttttatttga gacgagtctt gctctgtcgc taggctggag tgcagtggcg caatcttggc
29701 tcactgcaac ctccgactcc ttgttcaaag gattctcctg ccgcagcagc tagaattaca
29761 ggcacgtgcc accacgcccc gctaattttt tgtgttttta gtagagatag ggtttctcac
29821 catgtaggcc aggatggctc cgatctcctg acttcatgat tcgcccacct cgacctccaa
29881 aagtgtctgg attacaggcg tgagccattg cgctggcct atagaattct ttaaggcct
29941 aagttaaatg tgtgtgccta cagagaacat acgtatttaa atttgccagg ttgcagaggg
30001 cactacctac ttaacaaca ttacacgaaa ttcttagctt gagatttttg tattaccag
30061 gtagtatgaa ttcaggctgt aaactccctt gaggatcccc ttgaggatta tctaaaattt
30121 caggggagat tgtagttttc ctcttttagt cagtgttaag gtttgagaaa ggcatttttc
30181 ttgccatttc ctatggagtg gtacggtgtg ggtgaaggga gaggggctat ttccagttca
30241 acctgacact gacttttaag tcctttgggg tcccagctct agttggatgg tatattaaac
30301 tacatacctc ggatagacct tggacattgt ctccgtgtcc tgtgacctta taaactgtga
30361 aacaaaagct caagttcacc aagttccgca aatgcctca agttaaaact tgacttctgt
30421 ccaccttctt ttctgggttc ctactttcac atagtttttg tcttttgagt atttccaatt
30481 ctttttaagc tttggccaga agtttttagtt gtctgtagtt agagtgggtg tctagtgtac
30541 cataccactg aaacagaagc ctgttacttt tacaataata aaaacatatc tatgtgggct

```

FIGURE 8J

```

30601 ttttaaaaaa atttttaaat ttattttat ttattttttt ttgagatgga gtctcactct
30661 gtcgccagg ctggagtgc tggtcatgat cttggctcac tgcaacctct gccaccggg
30721 ttcaagcga tttcctgcct cagcctcctg agaagctgtg attacaggcg catgccaccg
30781 tgccctggcta aattttgtat ttttaggaga gacaggtttc accatgttgg tcaggctggg
30841 ctcaaactcc tgagctcagg tgatctgccc accttggcct cccaaagtgc tgggattaca
30901 agcatgagcc actgcacctg gccaaaccta tctatgtttt agttttttat gattatttta
30961 tattatccct tccccacaca tgaaataact tctttccaag tgacttatga agttgtcaac
31021 ttttcataaa gccttggaa aaagtgggca gaaaaattat aaataaaaag tcttgtctag
31081 gacagtacag gtctgtctta ttcctaaatt agatcagaat cgggttatgc cggtttttta
31141 taacatacca ttataattgg gtatgttaaa gaatgtatta gagatgcatt agaagagcga
31201 cctcattata agcctcttca cccatggatt ccaaggatat cttacaataa acctctggga
31261 taccttacgc tacagagcaa ctaaagtcca gctttagagc acaagggaat atgcaggata
31321 ttgggtcctg gaaactacaa aaaccatcaa actctacttt agggctaaaa ctttctttta
31381 atcaatctgt catatttatt gataaattag aaaatgtggc tgggtgcagt ggctcacacc
31441 tgtaatccta gcagtttggg aggccgaggt ggggtgatca cgagtgagac agatggagac
31501 cattctggct aacacagtga aaccctgtct tactaaaaat acaaaaatta gccgggctg
31561 gtgggtggcg cctgtagtcc cagctactca ggaggcggag gcaggagaat ggctgaacc
31621 cgggagacgg aggttgcagt gagccgagat tgcccactg cactccagcc tgggtgacag
31681 agggagactc catctcaaaa aaaaaaaaaa aaaagtacag gttcaaattt cattaattt
31741 tttttccatg ccccttctct ggtggtagga tgatagggga tgagtaaaat atcaaagaca
31801 actaaaaatt cctaaactag atcctcgggt cccaaagaag tgttaacctt ggagtgcacc
31861 accttctgag gttggtgctt cctggcacag gctcttgga cattctggat tgtctgacat
31921 tccttttact cgcttttaga cctaaagatg ttgctggggg aaaaggagc tagggagagg
31981 gaagggaag gatgtgggat aaggcataaa ctatgcttgt gggaaaaaaa caaccagtaa
32041 tttccttgat ggggtttttg cttgacttta aaatgcagta agcttttagg aagcttgatg
32101 gagectgtgt ttctaccatt agtctgcatt ttgtttggct taagatttca ttaagcttct
32161 taaaatggag atacttctgg aactcaaatt gctactgagc agggaatggc aaatcagtga
32221 actgtagga agtggaggga ttttgcaaa ctagagaaca caccttcac aggggcagcc
32281 acttctctgc tgggcccgct gttcatctgc agggacgtgc atcttggat tatttcaaa
32341 gtcagaactc agcattttta tgagaaatgt catgattttt aaggctttaa caagttatcc
32401 cacattaaaa aaataataat aaaccaggcc ggggtgcagt gctcacgcct gtaatccaa
32461 cattttggga ggctgagggt ggtggatcat gaggtcaggc gttcgagacc attatggcca
32521 acatggtgaa acactgtctc tacttaaaat acaaaaatta gctgggcatg gtggccggcg
32581 cctgtaatcc cagctactag ggaggctgag gcaggagaat cgcttgaacc tgggaggcgg
32641 aggaggtgc agtgaactga gatcgtaccc agcctgggca acagtgcag gttccgtctc
32701 aaaaaaaacc aaaaaacaca aaaaacaaaa acgacagaga aggccaaact aaacacactt
32761 gtgggctgga tgccgccatg cccaccgggt tgcgacctt gtgttggaact cttctgttca
32821 ccagacaccc tgccctgcga gaatgtatct catcctttgc tggagcaggt ttgcaggcac
32881 agtggagaga ggagagaaga aatgaaggga cacttatgca gaaccatgag tggccagaga
32941 ggaggagaag gaggtgaga ggagcaaaga agccatgaca acttcataat tctgagtga
33001 ctgggcagtg gccagaaatt ctgggtggtg atatgctgcc tttccaacag gtgaatatga
33061 aagaataagt caaacctgt tcaggacgct gttaattcca aatgtgaact ttttgagtca
33121 ttctttccat gtggaattca aaggagaatg taaacaaatt ttcaggagg acgtgcaata
33181 tccctgaaag ataacaaagt tcgtaacact tatttacata caacattctc tagttattga
33241 ttaacagat ctctacagac ttgcatgagg caacatttct taggcttgtt tgctacaata
33301 tctttaaaaa tacttgatta cacatcactt tagcttattt agatggactt ttcaccaagc
33361 tctgaactgg gatttcattt tgttgcatte atcctgctca cgagacacag gtaggcagca
33421 aatgagatta tccctccagt ccccatggat tggaaatgtt ccccttctt tatgagctca
33481 ctgcagtatc tccttctccc tttcccaaaa ggacagcctt tcctgcctca gggagaaga
33541 gagagacaga ctacagtgat ggagaccac tagatgtgca caagaggctg ccatccagt
33601 ctggagagga ccgagccgtg atgctgggtt ttgccatgat gggcttctca gtcctaattg
33661 tcttcttgc cggaacaacc attctaagc ctttatgct caggtaagaa acagaggaag
33721 gaaatctaga gtttccaatt cagataagat ccaggcacag cggtaaagga ggagaggctg
33781 gctttggtt ctcaatctgt gactctctgc catggtctag aaaaagaaaa atacaattcc
33841 tcttccgctg tcagtgtgg gggcagctgg gtgtggagga agaggtggcg agaaggtcaa
33901 agatatttcc ctgcactgcc tctcccac tcataatcta tatatactc caaacacgta
33961 atccacaaat tatagtttct tcathtaggt cataaatcct ccccttaaaa tgttggattt
34021 ccaagagaga gttaaaccct gtatgtgagc acaataaaag ttttttgagt ctgaattttt

```

FIGURE 8K

```

34081 tgagtttgac agtgtctacc tggcacatag tagttgctca atacatatta gtttccttcc
34141 ttttaattag gttctttatt caatatatgt agtgatacag ttgacctttg aacaacatgg
34201 gtttgaactt cgggaatcca cttataaatg gattttcttc tgtgcctgcc acccctgaga
34261 cagtaagatc aatccctcct ctttctcttc ctctcagcc tactcaacat gaagaggaca
34321 gggagaggac ctttatgatg atccacttcc atttagtaaa tagtaaacat gttttctctt
34381 ccttatgatt tttttcttc aatttttgtt ttaagttccg gggtagatgt acagaaatgtg
34441 caggtttgtt acaaaggtaa acgtgtgcc a tgggtggttg cagcacagat caaccatca
34501 cctgggtatt aagcccagca tgcattagct attcttcttg atgctctccc tccccctact
34561 ccactgagag gccccagtg gtgttgctcc cctctaggtg tccatgtgtt ctcatcttc
34621 agctcccact tttaagttag aacattcagt gtttggttct ctgtttctgc attagtttgc
34681 tgaggataat ggcttccagc ttcatttcat ctatgtcccc gcaaattgaca tgatcttgtt
34741 cttccttatg attttcttct cttttctttt ttctgagacg gagtcttgtc ctgtcaccca
34801 ggctgaagtg cagtggcacg atcttggttc actgcaacct ctgctcccg aataaagca
34861 gttctcctgc ctccagctcc caagtagctg ggaccacagg tgtgtgccac catacttggc
34921 taatttttaa attgttggtg gagatgggat ccccttatgt tgcctaggct ggtcgtgaac
34981 tcctaagctc aagtcactct tctaccttgg tctcccaaag tgctgggatt acaggcatga
35041 gccaccatgc ctgaccagta gttaaatttt tgaggagtca aaagtatatat gcaaattttc
35101 gactgcaggt ttgggggtgt ggtgttggtc tccctaaccct ctgaattgtt caaggttcaa
35161 ctttagttgc tacttattaa gtatatgtct tatgtcaat acatactata ttttactcta
35221 cttaaacctc cagcagccca tgatggaagt gttatttttag ttttcatttt ataaatgaga
35281 aagctgagtc tcagaaaagg caagtaactt gcccaagttc acatagccgg taaataatat
35341 gcctgtcatt ctgacgtaag acttgcttct taatacctaa atgatgtttg ctgaaatgag
35401 gggaatgggg gttctgatta gggaagggga ggggtctatt aattttgtag gatggccac
35461 acaacatgtg gtactctggg gccagggtcag ttcagggtaa aaggaaggca aggggtgctt
35521 tttggacatt gctttatttt tggacagccc tttttttttt ggacagccaa agagcaacag
35581 gaggccagg gggagtggaa gaggaaggcc taggagtcca gtgagggaaa ggggtcataa
35641 aaaaggaaga gagtggccta cttctgttat ttatgattca tctctgcca ttcctaggtt
35701 accattgtta ggggaaagat gtagaagta ccatttacta tagcacacag tccactgtt
35761 ggaagaattc cgcggatcct ctccctatag tgagtcgtat tagcggccgc aaatttatta
35821 gagcaatata gtccatacat gtcaagctcg accgatgccc ttgagagcct tcaaccagt
35881 cagctccttc cgggtggcgc ggggcatgac tattggcgcg ccggatcgat ccttaattaa
35941 gtctactaga taacttcgta taatgtatac tatacgaagt tattatctat gtcgggtgcg
36001 gagaaagagg taatgaaatg tgccccgctt acgcagggca tccattttatt actcaaccgt
36061 aaccgatttt gccaggttac gcggctgcag gcatgcaagc tttggcgcgc cgtcgaccaa
36121 ttctcatgtt tgacagctta tcatcgctta tctgccattc atccgcttat taccatttat
36181 tcaggcgtag caaccaggcg ttttaaggga ccaataactg ccttaaaaaa attacgcccc
36241 gccctgccac tcatcgagct actgttgtaa ttcattaagc attctgccga catggaagcc
36301 atcacaacac gcatgatgaa cctgaatcgc cagcggcatc agcaccttgt cgccttgctt
36361 ataatttttg cccatggtga aaacgggggc gaagaagtgt tccatatttg ccacgtttaa
36421 atcaaaactg gtgaaactca cccagggtat ggtgagacg aaaaacatat tctcaataaa
36481 cccttttagg aaataggcca ggttttacc gtaaacagcc acatcttgcg aatatatgtg
36541 tagaaaactgc cggaaatcgt cgtggtattc actccagagc gatgaaaacg tttcagtttg
36601 ctcatggaaa acggtgtaac aagggtgaac actatcccat atcaccagct caccgtcttt
36661 cattgccata cggaaactcg gatgagcatt catcaggcgg gcaagaatgt gaataaaggc
36721 cggataaaac ttgtgcttat ttttctttac ggtctttaaa aaggccgtaa tatccagctg
36781 aacggtcttg ttataggtac attgagcaac tgactgaaat gcctcaaaat gttctttacg
36841 atgccatttg gatatatcaa cgggtggtata tccagtgatt tttttctcca ttttagcttc
36901 cttagctcct gaaaatctcg ataactcaaa aaatacgcct ggtagtgate ttatttcatt
36961 atggtgaaag ttggaacctc ttacgtgccc atcaacgtct cattttcgcc aaaagttggc
37021 ccagggtctc cgggtatcaa cagggaacac aggtatttat tattctgcga agtgcatttc
37081 cgtcacaggt atttattcgc gataagctca tggagcggcg taaccgtcgc acaggaagga
37141 cagagaaagc gcggatcttg gaagtgcagg acagaacggg caggacctgg attggggagg
37201 cggttgcccgc cgctgctgct gacggtgtga cgttctctgt tccggtcaca ccacatacgt
37261 tccgcatctc ctatgcgatg cacatgctgt atgccggtat accgctgaaa gttctgcaaa
37321 gcctgatggg acataagtcc atcagttcaa cggaaagtcta cacgaaggtt tttgcgctgg
37381 atgtggctgc ccggcaccgg gtgcagtttg cgatgccgga gtctgatgag gttgcgatgc
37441 tgaaacaatt atcctgagaa taaatgcctt ggcctttata tggaaatgtg gaactgagtg
37501 gatatgctgt ttttgcctgt aaacagaga agctggctgt tatccactga gaagcgaacg

```

FIGURE 8L

```

37561 aaacagtcgg gaaaatctcc cattatcgta gagatccgca ttattaatct caggagcctg
37621 tgtagcgttt ataggaagta gtgttctgtc atgatgcctg caagcggtaa cgaaaacgat
37681 ttgaatatgc cttcaggaac aatagaaatc ttcgtgcggt gttacgttga agtggagcgg
37741 attatgtcag caatggacag aacaacctaa tgaacacaga accatgatgt ggtctgtcct
37801 tttacagcca gtagtgtctg ccgcagttga gcgacagggc gaagccctcg agtgagcgag
37861 gaagcaccag ggaacagcac ttatatattc tgcttacaca cgatgcctga aaaaacttcc
37921 cttggggtta tccacttacc caccgggata tttttataat tttttttttt atagttttta
37981 gatcttcttt ttttagagcg cttgtaggcc tttatccatg ctgggtctag agaaggtgtt
38041 gtgacaaatt gccctttcag tgtgacaaat caccctcaaa tgacagtcct gtctgtgaca
38101 aattgccctt aacctgtga caaattgccc tcagaagaag ctgttttttc acaaagttat
38161 ccctgcttat tgactctttt ttatttagtg tgacaatcta aaaacttgtc acacttcaca
38221 tggatctgtc atggcggaaa cagcgttat caatcacaa aaacgtaaaa atagcccgcg
38281 aatcgctccag tcaaacgacc tcaactgaggc ggcatatagt ctctcccggg atcaaaaacg
38341 tatgctgtat ctgttcgttg accagatcag aaaatctgat ggcaccctac aggaacatga
38401 cggtatctgc gagatccatg ttgtgaaata tgctgaaata ttcggattga cctctgcgga
38461 agccagtaag gatatacggc aggcattgaa gagtttcgcg gggaaggaag tggtttttta
38521 tcgccctgaa gaggatgccg gcgatgaaaa aggcattgaa tcttttccct ggtttatcaa
38581 acgtgcgcac agtccatcca gagggcttta cagtgtacat atcaacccat atctcattcc
38641 cttcttttatc ggtttacaga accggtttac gcagtttcgg cttagtgaag caaaagaaat
38701 caccaatccg tatgccatgc gtttatacga atccctgtgt cagtatcgta agccggatgg
38761 ctcaggcatc gtctctctga aaatcgactg gatcatagag cgttaccagc tgctctaaag
38821 ttaccagcgt atgctctgact tccgcgcgcg ctctctgcag gtctgtgtta atgagatcaa
38881 cagcagaact ccaatgcgcc tctcatatcat tgagaaaaag aaagggcgcc agacgactca
38941 tatcgatatt tccttccgcg atatcacttc catgacgaca ggatagtctg agggttatct
39001 gtcacagatt tgagggtggt tcgtcacatt tgttctgacc tactgagggt aatttgtcac
39061 agttttgctg tttccttcag cctgcatgga ttttctcata ctttttgaa tgtaattttt
39121 aaggaagcca aatttgaggg cagtttgtca cagttgattt ccttctcttt cccttcgtca
39181 tgtgacctga tatcgggggg tagttcgta tcattgatga gggttgatta tcacagttta
39241 ttactctgaa ttggctatcc gcgtgtgtac ctctacctgg agtttttccc acggtggata
39301 tttcttcttg cgtctgagcg aagagctatc tgacagaaca gttctctttt gttcctcgc
39361 cagttcgctc gctatgctcg gttacacggc tgccggcgagc gctagtata ataagtgact
39421 gaggtatgtg ctcttcttat ctcttttgt agtgttgctc ttatttttaa caactttgcg
39481 gttttttgat gactttgcga tttgtgtgtt gctttgcagt aaattgcaag atttaataaa
39541 aaaacgcaaa gcaatgatta aaggatgttc agaattgaaac tcatggaaac acttaaccag
39601 tgcataaacg ctggtcatga aatgacgaag gctatcgcca ttgcacagtt taatgatgac
39661 agcccggaag cgaggaaaat aaccggcgcg tggagaatag gtgagcagc ggattttagt
39721 ggggtttctt ctccaggtat cagagatgcc gagaaaagcag ggcgactacc gcaccggat
39781 atggaaattc gaggacgggt tgagcaacgt gttggttata caattgaaca aattaatcat
39841 atgcgtgatg tgtttggtac gcgattgcga cgtgctgaag acgtatttcc accggtgatc
39901 ggggttgctg ccataaaagg tggcggttac aaaacctcag tttctgttca tcttgctcag
39961 gatctggctc tgaaggggct acgtgttttg ctctggaag gtaacgaccc ccagggaaca
40021 gcctcaatgt atcacggatg ggtaccagat ctctcatattc atgcagaaga cactctcctg
40081 cctttctatc ttggggaaaa ggacgatgtc acttatgcaa taaagcccac ttgctggccg
40141 gggcttgaca ttattccttc ctgtctggct ctgcaccgta ttgaaactga gttaatgggc
40201 aaatttgatg aaggtaaaact gccaccgat ccacacctga tgctccgact ggccattgaa
40261 actgttgctc atgactatga tgtcatagtt attgacagcg cgcctaacct gggatatcggc
40321 acgattaatg tcgtatgtgc tgctgatgtg ctgattgttc ccacgcctgc tgagttgttt
40381 gactacacct ccgcaactga gtttttcgat atgcttcgtg atctgctcaa gaacgttgat
40441 cttaaagggt tcgagcctga tgtacgnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnngac
40501 agtctatctg ttagttatat tctagccctt tatagcttgc tggctctcaag atgtagaatt
40561 gctgaaagat gtaattcaga ctgtttggaa agaagacaag tcctccctcc ttcttgggtt
40621 cactgtgaac caccgaggtt tctacgatac atgggtctaac cactgaaact ttaacttctc
40681 tactgagata ttttagaagt gtaaaataca gactgtattt caaagacagc atgaaaagc
40741 cagcttttgt agcaggcacc aagcttggtt tactttatac aggcctcccta attcctctct
40801 gccacttcaa ccttcaactg atatccctgg cagaaggatg aaaaatcaaa agggcaagag gattcatttc
40861 atagaacatt tgtttcaggt attgggacaa ctccaagtat taagataaca gattcatttc
40921 atgctgtttc atttaataca tgtttattca tggcaaat ttttttttat tttttttatt
40981 ttttttttta atttaagtgg gtacatagta gttttatgta tttatgggtt acgtgagata

```

FIGURE 8M

```

41041 ttttgataca ggcatgcaat aatcacatca gggtaaatgg ggtgtccatc atctcaagta
41101 tttatccttg tgtcacaaac aatccaatta tactcttagt tgtttaaaaa tgcggggagtg
41161 ggaggggcaag tggaggggaga gcattaggac aaatacctaa tgcattgcggg gcttaaaacc
41221 tatgtgatgg gttgacaggt gcagcaaacc accatggcac atgtatacct atgtaacaaa
41281 cctgcacggt ctgcacatgt atcccgtaat gtaaaagtaaa ataaaaatagc ataaaaatcaa
41341 ataaaggaat cactaatgaa acctacattt gcaaaataaa ttataaatat tgaaaagaaa
41401 taaaaagtac aaaaaattat ttttgtctat agtcactcta ttgtgctatc aaatactagt
41461 cttatttcatt ctttttgttt tttgtaccca ctaaccatcc cactccccgc cctacacccc
41521 ccaacaaccc gccccatact cttccaagcc tctggtaacc gtccttctac tatctccatg
41581 agtcgaattg ttttaatttt taacttccac aaataagtga gaatatgtga agtttgtctt
41641 tctgtacctg gcttaaatat cacttaacat aatgacctcc agttccatcc atgttattgc
41701 aaatgactgg acctcatttt cttaatgggt gaatagtact ccattgtgta tatgtaccac
41761 attaaaaaaa atcaatttgt tagttgatgg attgctttca aatcttgggt attgccaata
41821 gtgctgcaat agctaagatt tggatatctt tgggagtgtg gatattcttt ggatatcttt
41881 taatcttttc ttttctttt gagacagagt ctgcgtctgt tgcccagggt ggagtgtagt
41941 ggcgcaatct tggctcactg caacctctgc ctcccgggtt caaacaattc tctgtcttca
42001 gcctcctgag tagcttggat tacaggcact caccaccatg cccagataat ttttgtattt
42061 ttagtagaga cggggtttca ccatgttggc caggctgggtc tcaaactcct gacctcagat
42121 gatccaccca tctcgccctc ccaaaagtgt gggattacag gcgtgagcca cctcgcccag
42181 cctgattttt tttcttttga gtatatacct acctctattt tcagtttttt gaggcacctc
42241 caaactgttc tccatagtga ttgtactaat ttacattccc actaacagtg tacaaggggt
42301 cctttttctc cctattcttg ctgacatttg ttattgcttg acttttggat aaaagccatt
42361 ttaactgggg tgagatgata tgtcattgta gttttgattt gcatttctct aatgatcaat
42421 gacaatgcct ttttgtatgc ttgtttccca cttgtatgtc ttcctttgag aaatgtctat
42481 gcagatcttt tgcccattta aaaaactgga ttattatatt ttttctata gagttgtttg
42541 agctcctttt atattctgat tattaacccc ttgtcagatg ggtagtttgc aattattttc
42601 taccactctg tggtttgtct cttcactttg ttgattcttt cttttgctat gcagaaagtt
42661 ttttaatttga tatgatccta tgtgttcaca tttgctttgg ttgctgtgtc ttatgaggtt
42721 ttttcaagg aatctttgcc tactctggga cttactgttt tcttttacta gtttcatagt
42781 tctgaggtctt aaatttcagt ctttaatacca ttttgatttg atttttgtat aaggatagag
42841 ataacaggtc tagttcactc ttttgcataat ggatatccag ttttcccagc atcatattat
42901 gaagagactg tctgtttccc aatttatgct cttggttaact ttgtcaaaaa tgaattcact
42961 gcagatgtat gcatttacct ctgggttctc tattctgttc cactgatgta tgtgtctgtt
43021 tttatgccag taccttgtctg ttttggttac tgtagtttta tagtataagt ttaagtcaga
43081 taaagtgatt tctcccattt tgttcttttt gctcagggtg gctttgggtc ttctgggtct
43141 tctgtgggtc caaatgaatt ttagaataatt ttttctattt ctgtgaagaa tgtcttgggt
43201 attttgatag gcattgtcatt aaatctgtag atcgctttga gtaatatgga cattttaaca
43261 ttattgatat ttccaatcca tgaacatgta atgtctttcc ttttttgtgg cttcttcaat
43321 ttcttgcatt aatgttttat agttttcatt ttagatatca ttcacttcgt tagttaattc
43381 ctaggatatt aattttattt gtatgtattg taaatgggat cttttaaaat ttctttttca
43441 gattgttcac tgttagcata tagaaatgct actgatgttt gtatgttgaa tttgtatcct
43501 gcaactttac tgagtttgtt tatcagttct tatagttttc ttgatgggat ctttagtatt
43561 tcccaaatat aagatcatat ctgaaaacta gaataatttg acttcttctc atocaaattg
43621 gatgcccttt atttctttct cttgtatgat tgctcaagct aggactttca gtattatgtt
43681 gaataacagt ggtgtaagtg ggcatccttg ttgttttcca gatcttagag aaaaggcttt
43741 caatttttcc ccattcagta tgatactaac tctggatctg ttgtatatgc tttttattat
43801 gttgaggtat gttccttctg taaccagttt tgtgaggttt ttgttatgaa gggctgttga
43861 attttatcaa gtgctttttc agcatcaatt gaaatgatca tatagttttg tctttcatte
43921 tgttgatatg atgtattaca ttaattgatt tgcatatgtt gaactatctt tgcattccca
43981 ggctaaatct cacttcgtca taatgaatga tgagtgaatt agttagtga tttgatgag
44041 ttatcgaata ttgagtttga tttggtagta tttgagaatt tttgcatcaa tttcatcag
44101 aaatattggt gtatagtttt atttttattg atgtgtcttt gtctagctct agtataaga
44161 taatactggc ctcatagaat gagtctggaa gtattccctc ctttctttgt gtttttgaat
44221 agtttgagta ggattgggtat agtttttttt tatttaattt gtttatttgt cttgagacac
44281 ggtgtcactc tgtcaccaca ggctggagta cagtgggtgt atcttggctt accatagcct
44341 ctgcctcgcc tcccaggctc aagcgattct ctaataatat ttgctttata tatctgggtg
44401 ctccagtgtt tgggtgcata atatttatca aattgttata tcctcttgcct gaattgacct
44461 ctttatcatt atatagtgc cttcttgttc tcttcttata gtttttgcct tgaaatctat

```

FIGURE 8N

```

44521 ttgtctgatt tatccaattt atttatctga ctctgctctt ttttttgggt tccatttgca
44581 tggaatattt ttttccatcc cttattttca gtttactgtg tattcttggg caagttactt
44641 tctgagcctc aatttcctca tctgatagtt ggtgtctgtt aggatctatc ttacaggact
44701 gttttaagga ttaaatagaga caatgcacag tgcctggggc atgatgagat atggtaatta
44761 ggaaataatt actcaggcca ggcacgggtg ctcatgccta taattccaac attttgggag
44821 gctatggtgg gaggatcgct taacaccagc ctgggcaata tagtcacacc cagtctctcc
44881 aaagaaagaa agaaagaaag aaaaaaatta gccactatt gtggtcacat gactgtagtc
44941 ccagctactc aggaggctga gatgggagga tcacttgagc ctggaagggt gagggtgagc
45001 tgagccatga ttgtgctact gcgctccagc ttgggcaaca acatgagaac ttatttcaaa
45061 atagctaaca aatggtagtt attattgtta catttattta tttagaatta ttattcaatt
45121 attattatgt aataatcaac aaaaattatc cttttttgta gcatcttttt atctcttctc
45181 tgatttcttt tttgaacaaa tttaaaaaca tattataata ttaaaaaacg gagaccagggt
45241 ctccctgtgt taccaggct gtcctcaaac tcctgggctc gagcaatcct tccgccacag
45301 ctttccaaaag tgctgcggtt acaggcgtga gccactgtgc ctggccttgc attatttctg
45361 ttatgggttat ttgtttattt gttatctgtg taatttattt actttgatgt ttaaagcaaa
45421 gaattacaac tcagtgaagg agatatacctc acatgtaact tgcaactcta aatattgacct
45481 tatctaagag ggtattagtt tcaattatgc tgccctatg ggtcattgaa gtgagactct
45541 tctgtgtgtc ccataagagt gcactaaggc aactgaagtc atttgatgtc ctcagccccc
45601 ttggcaagca cagagggtgt caaaagactg atgaacttgg cactaagatt tttcttccag
45661 aagacaggga gaaatcacca aatggttttt tttttaagc aaactaaata tttagccacc
45721 ttttgggttc actttttttt ttcttttctt tttttactag tcattgactt gaagctcatg
45781 gttttactct ttaacaccag ccagtcactc gcaggttctt gggacaaaaa gcaaaaggaa
45841 taatgatccc ttgtggattc ctgagtcacc tgctttgtgg gctgtgcagc gctgtgtgtc
45901 agcaccacac cggctacctc actattgtct agggtttgtg ctttggggagc tagcctggaa
45961 gctaagaact ctcactttca cctgggtatta actacttgtt ttcttttttt tttttttttt
46021 tgagacggag tctcgctctt gtcacccagg ctggagtgtc gtggcacgat ctcagctcac
46081 tgcaacctct gcctcctggg ttcaagcgat tctcctgcct cagcctcccg agtagctggg
46141 attaaaggct cctgccacca ccccgagcta atttttttta tttttagtag agatggagtt
46201 tcattcatgtt gggcaggctg gcctcgaaact tctgacctcc ggtgatcctc ccaccttggc
46261 ctctgaagt gctgggatta caggcgtgag ccacattgcc cagcctgtga ttaactactt
46321 tctaaaggcg cttcctgttg cacaagaatg ctgtttgcag gaatgcctgt agacacatga
46381 accctgggcc tgctgtctcc ttaatttgag ggaagctatg gatgtcaagc agaatagagc
46441 tattttgtta cttgaccagg ttccataaac tttttgcaca tacctttatt cctgcacctg
46501 tcccacataa ttatcccttt atagctcttt caccagcagg aatataaatg tcatggcagg
46561 gccaccatgt tgcaaaactc ctagggtcca tcctgcagtt actagcttgt gtgaatggta
46621 cttcttgag gtaccattgc aatgcacagc ctgagcagcc ctctgcagta agtccaactg
46681 tggatggggc aaaatgagtg gttagcagaa ttggcctcat acatctttta ttctgtcatt
46741 gagtatgtgg taggcactca agaaatattt ttaaataaat gacattggtc ttccatttta
46801 ttttgagtta atttgacca gcttaatcag tggatctttt caactagtta attgtcctac
46861 ccttccaccc aggaagggtt ttcagtttta atagattttt gttattgggt tttttttgca
46921 cattcagcaa aacctctgga aagaggagtt catatcagcc cctcatgcag agggagctgt
46981 gagtattcag gctgcttggg gctgctgcgg tctttccagc agaggtgggt ggtcctagag
47041 tgggtggttg gtccattctt gacctatgat gatgctggga aataactcag tgatgtgggtg
47101 tcattttttt atctcacaga tgtggctctg gagaagagat ttaagtatat atcttcagaa
47161 catagcatca atgagtttgg gtcataagct cagccaactc agccagaaac agatacacag
47221 tatgcaacca agtggaata aattcaaccc attttcctaa tttggaatat agagcctgag
47281 ggggaagtt gtgggtggat tatctgtcct gggttcttcc tatttcttga atttatactt
47341 tagtgaaggg aaattttgct aactgttaat gtgccagaat gggaaagccc cctacctagg
47401 ccctcacctc tgaatttttc cagttcaagt ggacaagtag tttttgagca ctcactatgt
47461 gtcaggctca ggggtgacct tgcgatttcc acttctctct gatcttttca cctaagcctg
47521 tgggacttgt gttctgggag aaaattagaa gttacatcat ccttatgggt ctgaaaaaat
47581 tgagctcccc agacttgggt tcgcctgata tgcttctcaa atgttaaaga ggaacataaa
47641 gccatgtaga aaatcctttt cttttgacaa agcctggctt tcaaattcta ctgttgctgc
47701 agtcacaaa attatacatt tgaatgtcag agattgaatg ttgacatcct tattgcaaaa
47761 atcctatccc cacttctggc ttagctgccc cagagagctc tgtgtctcgg gctgactaga
47821 gaaaagacca cacagttaga gatgcaaaaa aggaaatgcc attaacatat ttcagggtga
47881 atatttcttt gcatttggca ctttgcaatc tagtccaggc ttaagatttg taggatttct
47941 tttgctgttt cccttttttt acttgtcagg gaaacaaggc ataaatattt cagcttggac

```

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/19585

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :Please See Extra Sheet.

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1, 23.5, 24.3, 24.31; 530/350, 387.1, 387.9; 435/ 69.1, 71.1, 71.2, 471, 320.1, 254.3, 254.11, 325, 6, 7.1, 7.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
~~NONE~~

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,637,470 A (KACZOROWSKI et al.) 10 June 1997 (10.06.97), see entire document.	1-42
A	US 5,776,734 A (KACZOROWSKI et al.) 07 July 1998 (07.07.98), see entire document.	1-42

☐

Further documents are listed in the continuation of Box C.

☐

See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

20 NOVEMBER 2000

Date of mailing of the international search report

27 DEC 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PREMA MERTZ

Telephone No. (703) 305-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/19585

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

C07K 14/47, 16/18; G01N 33/53, 33/567; C12N 5/10, 15/12, 15/63, 15/64

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

536/23.1, 23.5, 24.3, 24.31; 530/350, 387.1, 387.9; 435/ 69.1, 71.1, 71.2, 471, 320.1, 254.3, 254.11, 325, 6, 7.1, 7.2

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

WEST, CAS ONLINE, MEDLINE, CAPLUS

search terms: calcium sensitive potassium channel beta2, beta2a, beta2b, beta2c, beta2d, nucleic acid, recombinant production, antibody, assay, binding, activator, inhibitor, gene transcription, nuclear factors